



BlueCross BlueShield
of Alabama

Name of Policy:

Continuous or Intermittent Monitoring of Glucose in the Interstitial Fluid

Policy #: 038
Category: DME

Latest Review Date: May 2018
Policy Grade: B

Background/Definitions:

As a general rule, benefits are payable under Blue Cross and Blue Shield of Alabama health plans only in cases of medical necessity and only if services or supplies are not investigational, provided the customer group contracts have such coverage.

The following Association Technology Evaluation Criteria must be met for a service/supply to be considered for coverage:

- 1. The technology must have final approval from the appropriate government regulatory bodies;*
- 2. The scientific evidence must permit conclusions concerning the effect of the technology on health outcomes;*
- 3. The technology must improve the net health outcome;*
- 4. The technology must be as beneficial as any established alternatives;*
- 5. The improvement must be attainable outside the investigational setting.*

Medical Necessity means that health care services (e.g., procedures, treatments, supplies, devices, equipment, facilities or drugs) that a physician, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury or disease or its symptoms, and that are:

- 1. In accordance with generally accepted standards of medical practice; and*
- 2. Clinically appropriate in terms of type, frequency, extent, site and duration and considered effective for the patient's illness, injury or disease; and*
- 3. Not primarily for the convenience of the patient, physician or other health care provider; and*
- 4. Not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.*

Description of Procedure or Service:

Tight glucose control in patients with diabetes has been associated with improved outcomes. Several devices are available to measure glucose levels automatically and frequently (e.g., every 5-10 minutes). The devices measure glucose in the interstitial fluid and are approved as adjuncts to traditional self-monitoring of blood glucose levels. Devices can be used on an intermittent (short-term) basis or a continuous (long-term) basis.

Hypoglycemia affects many aspects of cognitive function, including attention, memory, and psychomotor and spatial ability. Severe hypoglycemia can cause serious morbidity affecting the central nervous system (e.g., coma, seizure, transient ischemic attack, stroke), heart (e.g., cardiac arrhythmia, myocardial ischemia, infarction), eye (e.g., vitreous hemorrhage, worsening of retinopathy), as well as cause hypothermia and accidents that may lead to injury. Fear of hypoglycemia symptoms can also cause decreased motivation to adhere strictly to intensive insulin treatment regimens.

The advent of blood glucose monitors for use by patients in the home revolutionized the management of diabetes. Using fingersticks, patients could monitor their blood glucose level both to determine the adequacy of hyperglycemia control and to evaluate hypoglycemic episodes. Randomized controlled trials of tight control have demonstrated benefits for type 1 diabetics in decreasing microvascular complications. The impact of tight control on type 1 diabetic patients and on macrovascular complications such as stroke or myocardial infarction is less certain. The Diabetes Control and Complications Trial (2002) demonstrated that a relative HbA1c level reduction of 10% is clinically meaningful and corresponds to approximately a 40% decrease in risk for progression of diabetic retinopathy and 25% decrease in risk for progression of renal disease.

Due to an increase in turnover of red blood cells during pregnancy, HbA1c is slightly lower in women with a normal pregnancy compared with nonpregnant women. The target A1C in women with diabetes is also lower in pregnancy. The American Diabetes Association recommends that, if achievable without significant hypoglycemia, the A1c should range between 6.0 to 6.5%; an A1c less than 6% may be optimal as the pregnancy progresses.

Tight glucose control requires multiple daily measurements of blood glucose each day (i.e., before meals and at bedtime), a commitment that some patients may be unwilling or unable to meet. In addition, the goal of tight glucose control has to be balanced with an associated risk of hypoglycemia. Hypoglycemia is known to be a risk in patients with type 1 diabetes. Studies have also found that approximately 50% of patients with type 2 diabetes may experience hypoglycemic episodes, but the severity of these episodes may vary. An additional limitation of periodic self-measurements of blood glucose is that glucose values are seen in isolation, and trends in glucose levels are undetected. For example, while a diabetic's fasting blood glucose level might be within normal values, hyperglycemia might be undetected postprandially, leading to elevated hemoglobin A1C values.

Recently, measurements of glucose in interstitial fluid have been developed as a technique of automatically measuring glucose values throughout the day, producing data that show the trends in glucose measurements, in contrast to the isolated glucose measurements of the traditional

blood glucose measurements. Although devices measure glucose in interstitial fluid on a periodic rather than a continuous basis, this type of monitoring is referred to as continuous glucose monitoring (CGM).

Several devices have received U.S. Food and Drug Administration (FDA) approval. The first 2 approved devices were the Continuous Glucose Monitoring System (CGMS[®]) (MiniMed), which uses an implanted temporary sensor in the subcutaneous tissues, and the GlucoWatch G2[®] Biographer, an external device worn like a wristwatch that measures glucose in interstitial fluid extracted through the skin with an electric current (referred to as reverse iontophoresis).

Devices that have subsequently been approved include those for pediatric use and those with more advanced software, more frequent measurements of glucose levels, more sophisticated alarm systems, etc. Devices initially measured interstitial glucose every 5 to 10 minutes and stored data for download and retrospective evaluation by a clinician. With currently available devices, the time intervals at which interstitial glucose is measured ranges from every 1 to 2 minutes to 5 minutes and most provide measurements in real-time directly to patients. While continuous glucose monitors potentially eliminate or decrease the number of required daily fingersticks, it should be noted that, according to the FDA labeling, monitors are not intended to be an alternative to traditional self-monitoring of blood glucose levels but rather provide adjunct monitoring, supplying additional information on glucose trends that are not available from self-monitoring. In addition, it is important to note that devices may be used intermittently, e.g., time periods of 72 hours, or continuously, i.e., on a long-term basis.

In addition to stand-alone continuous glucose monitors, several insulin pump systems have a built-in CGM. This evidence review addresses CGM devices, not the insulin pump portion of these systems.

For coverage information regarding Medtronic's MiniMed 530G/630G and 670G, refer to medical policy #636 Artificial Pancreas Device Systems.

Policy:

Effective for dates of service on or after September 26, 2013:

Intermittent Monitoring

Intermittent monitoring, i.e., 72 hours, of glucose levels in interstitial fluid meets Blue Cross and Blue Shield of Alabama's medical criteria for coverage in patients with **Type 1 diabetes mellitus whose diabetes is documented in the medical records as *poorly controlled despite current use of **best practices.**

*Poorly controlled Type 1 diabetes mellitus includes the following clinical situations:

- Unexplained hypoglycemic episodes;
- Hypoglycemic unawareness;
- Suspected postprandial hyperglycemia;
- Recurrent diabetic ketoacidosis.

Intermittent monitoring of glucose levels in interstitial meets Blue Cross and Blue Shield of Alabama's medical criteria for coverage in patients with Type 1 diabetes prior to insulin pump initiation to determine basal insulin levels.

Intermittent monitoring is generally conducted in 72-hour periods. It may be repeated at a subsequent time depending on the patient's level of diabetes control.

Continuous Monitoring

Continuous, i.e., long-term, monitoring of glucose levels in interstitial fluid, including real-time monitoring, as a technique of diabetic monitoring, meets Blue Cross and Blue Shield of Alabama's medical criteria for coverage when the following situations are documented in the medical records and occur despite use of **best practices:

- Patients with Type I diabetes who have recurrent, unexplained, severe (generally blood glucose levels less than 50 mg/dL) hypoglycemia or impaired awareness for whom hypoglycemia puts the patient or others at risk; or
- Patients with Type I diabetes who are pregnant.

** Best practices in diabetes control for patients with diabetes mellitus include:

- Compliance with a regimen of four (4) or more fingersticks each day; or
- Use of insulin pump; or
- Prior use of intermittent (72-hour) glucose.

Replacement or upgrade of existing, properly functioning equipment, even if warranty has expired, does not meet Blue Cross and Blue Shield of Alabama's medical criteria for coverage.

The **glucose sensor and transmitter components** of a continuous glucose monitor used with a combined continuous subcutaneous insulin infusion and blood glucose monitoring devices **meet** Blue Cross and Blue Shield of Alabama's medical criteria for coverage when all the above criteria met.

Other uses of continuous monitoring of glucose levels in interstitial fluid as a technique of diabetic monitoring do not meet Blue Cross and Blue Shield of Alabama's medical criteria for coverage and are considered **investigational.**

Coverage for non-medical items, even when the items may be used to serve a medical purpose, such as smart devices (smart phones, tablets, personal computers, etc.) are non-covered. This includes smart devices used in conjunction with Continuous Glucose Monitors.

For coverage information regarding Medtronic's MiniMed 530G/630G and 670G, refer to medical policy #636 Artificial Pancreas Device Systems.

Blue Cross and Blue Shield of Alabama does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Cross and Blue Shield of Alabama administers benefits based on the member's contract and

corporate medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.

Key Points:

The most recent literature search was performed through January 08, 2018. Following is a summary of the key literature to date:

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function-including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Most of the following discussion focuses on the clinical utility of continuous glucose monitoring (CGM) systems. That is, their ability to provide either additional information on glucose levels, leading to improved glucose control or to improve the morbidity/mortality associated with clinically significant severe and acute hypoglycemic or hyperglycemic events. Because diabetic control encompasses numerous variables including the diabetic regimen and patient self-management, randomized controlled trials (RCTs) are important to isolate the contribution of interstitial glucose measurements to the overall diabetic management. Data on patients with Type I diabetes and Type II diabetes are discussed separately.

Type I Diabetes

This evidence review includes RCTs that report on outcomes of CGM devices. CGM devices will be categorized as continuous, long term, monitoring devices by the patient to direct insulin regimens, and intermittent (i.e., 72 hour) periodic monitoring used by the provider to optimize management.

CGM Devices for Continuous, Long Term Use

Systematic Reviews

A number of systematic reviews and meta-analyses of RCTs evaluating CGM for long-term, daily use in treating type 1 diabetes have been published. These systematic reviews have focused on slightly different populations, and some did not separate long-term CGM from intermittent glucose monitoring. The most recent meta-analysis, which was also the only analysis that used individual patient data, was published by Benkhadra et al in 2017. The meta-analysis evaluated data from 11 RCTs that enrolled patients with type 1 diabetes and compared real-time CGM to a control intervention. Studies in which patients used insulin pumps or received multiple daily insulin injections were included. Reviewers contacted corresponding study authors requesting individual patient data; data were not obtained for 1 trial. Mean baseline hemoglobin A1C (HbA1c) levels were 8.2% in adults and 8.3% in children and adolescents. The overall risk of bias in the studies was judged to be moderate. In pooled analyses, there was a statistically significantly greater decrease in HbA1c levels with real-time CGM versus control conditions. Overall, the degree of difference between groups was 0.26%. In subgroup analyses by age, there was significantly greater change in HbA1c levels among individuals 15 years and older, but not in the younger age groups. There were no significant differences between groups in the time spent in hypoglycemia or in the incidence of hypoglycemic events. Key findings are shown in Table 2.

Table 2. Main Findings from a 2017 Individual Patient Data Meta-Analysis on Real-Time CGM in Type 1 Diabetes

No of Trials	N	Outcomes	Point Value	95% Confidence Interval	p
Change in HbA1C levels, %					
8	1371	Overall	-0.258	0.464 to -0.052	0.014
7	902	Age > 15 y	-0.356	0.551 to -0.160	<0.001
7	178	Age 13-15 y	-0.039	-0.320 to 0.242	0.787
7	291	Age ≤ 12 y	-0.047	0.217 to 0.124	0.592
Time Spent in Hypoglycemia <60mg/dl, min					
4	706	Overall	-8.549	-31.083 to 13.985	0.457
4	467	Age > 15 y	-8.095	-32.615 to 16.425	0.518
3	109	Age 13-15	-13.966	31.782 to 3.852	0.124
3	130	Age ≤ 12 y	-9.366	19.898-1.167	0.081
Incidence of Hypoglycemic events <70mg/dl, mean no. events					
3	351	Overall	0.051	-0.314 to 0.416	0.785
3	277	Age >15 y	-0.074	-0.517-0.368	0.742
2	47	Age 13-15 y	0.536	0.243 to 1.316	0.177
2	27	Age ≤ 12 y	0.392	0.070 to 0.854	0.097

CGM: continuous glucose monitoring; HbA1C, hemoglobin A1C

Adapted from Benkhadra et al (2017)

Earlier meta-analyses of glucose monitoring devices for type 1 diabetes tended to combine studies of intermittent glucose monitoring with studies of long-term CGM. Several reported separate subgroup analyses for long-term CGM. A 2012 Cochrane review of CGM in type 1 diabetes in adults and children included RCTs comparing CGM with conventional self-monitored blood glucose (SMBG). In pooled analysis (6 studies; n=963 patients) of studies of long-term CGM, the average decline in HbA1c levels 6 months after baseline was statistically significantly larger for CGM users than for SMBG users (mean difference [MD] change, -0.2%; 95% confidence interval [CI], -0.4% to -0.1%), but there was no difference in decline in HbA1c levels at 12 months (1 study, n=154 patients; MD change, 0.1; 95% CI, -0.5 to 0.7). In a meta-analysis of 4 RCTs (n=689 patients), there was no significant difference in the risk of severe hypoglycemia between CGM and SMBG users and the confidence interval for the relative risk (RR) was wide (RR=1.05; 95% CI, 0.63 to 1.77), indicating lack of precision in estimating the effect of CGM on hypoglycemia risk. Reviewers were unable to compare longer term change in HbA1c levels or hypoglycemia outcomes for real-time CGM. Trials reporting results by compliance subgroups found larger treatment effects in highly compliant patients.

A 2011 systemic review of RCTs on CGM included trials conducted in adults and children with Type I diabetes who were on an intensive insulin regimen (studies of Type II diabetes were not included). This meta-analysis required a minimum of 12 weeks of follow-up in the studies (as compared to at least 8 weeks in the Gandhi meta-analysis). Studies compared CGM to SMBG; there was no restriction related to type of CGM device, but the CGM readings had to be used to adjust insulin dose or modify diet. A total of 14 RCTs met eligibility criteria. Study duration ranged from 3 to 6 months. The baseline mean HbA1c ranged from 6.4 to 10. Five of the included studies found a statistically significant decrease in HbA1c in favor of CGM while 9 did not. In a pooled analysis, there was a statistically significant reduction in HbA1c with CGM compared to SMBG, WMD: -0.26%, 95% CI: -0.34 to -0.19%. For the subgroup of 7 studies that reported on continuous long term monitoring, this difference was statistically significant (WMD = -0.26; 95% CI, -0.34, -0.18). In a sub-group analysis by age, there were significant reductions in HbA1c with CGM in studies of adults (n=5), WMD: -0.33 (95% CI: -0.46 to -0.20) and in studies with children and/or adolescents (n=8), WMD: -0.25, 95% CI: -0.43 to -0.08. Four of the studies provided data on the frequency of hypoglycemic episodes. Pooled results showed a significant reduction in hypoglycemic events in CGM versus SMBG (SMD=-0.32; 95% CI,-0.52 to -0.13). Five of the studies reported the percentage of patients with severe hypoglycemic episodes and there were no differences in the percentage of patients with severe hypoglycemic episodes for CGM versus SMBG in any of the 5 studies.

Randomized Controlled Trials

Recent RCTs not included in the meta-analyses are described next.

Van Beers et al (2016) published a crossover RCT comparing CGM with SMBG and focusing on patients with impaired hypoglycemia awareness. Eligible patients were 18 to 75 years old, were treated with insulin infusion pumps or multiple daily insulin injections, undertook at least 3 SMBG measurements per day, and had impaired awareness of hypoglycemia (i.e., Gold score ≥ 413). The trial used an artificial pancreas device system without using the low glucose suspend feature. After a 6-week run-in phase (during which patients received education about diabetes management), 52 patients received both 16 weeks of CGM and 16 weeks of SMBG, in random

order. There was a 12-week washout period between interventions. All patients were included in the primary intention-to-treat analysis. Six patients withdrew from the study early.

The primary outcome, time spent in normoglycemia (4-10 mmol/L), was significantly higher in the CGM phase than in the SMBG phase. The percentage of time spent in normoglycemia was 65.0% in the CGM phase and 55.4% in the SMBG group (mean difference, 9.6%; $p < 0.001$). The sequence allocation did not have an effect on the primary end point. Most other CGM-derived outcomes (e.g., number and duration of nocturnal hypoglycemia events) also significantly favored the CGM group. The total number of severe hypoglycemic events (i.e., those needing third-party assistance) was 14 in the CGM phase and 34 in the SMBG phase, which differed significantly between groups ($p = 0.033$). The number of patients with 1 or more severe hypoglycemic event during the intervention period, however, did not differ significantly between phases (10 in the CGM phase and 18 in the SMBG phase ($p = 0.062$)). HbA1c outcomes did not differ significantly; e.g., change in HbA1c levels from baseline was -0.1% in both phases ($p = 0.449$). In terms of hypoglycemia awareness, one of 4 variables, Gold score at the study end point differed significantly (mean, 4.6 for the CGM phase vs 5.0 for the SMBG phase, $p = 0.035$.); 3 other variables related to hypoglycemia awareness did not differ between groups.

Two 2017 RCTs evaluated long-term CGM in patients with type 1 diabetes treated with multiple daily insulin injections. Both trials used the Dexcom G4 CGM device. Lind et al (2017) reported on a crossover study with 142 adults ages 18 and older who had baseline HbA1c levels of 7.5% or higher (mean baseline HbA1c level, approximately 8.5%). There was a 6-week run-in period using a CGM device with masked data and patients were excluded from further participation if they did not believe they would use the device more than 80% of the time or did not perform an adequate number of calibrations during the run-in period. Enrolled patients underwent 26-week treatment periods with a CGM device and conventional therapy using SMBG, in random order. There was a 17-week washout period between intervention phases. The primary end point was the difference in HbA1c levels at the end of each treatment period. Mean HbA1c levels were 7.9% during CGM use and 8.4% during conventional therapy (mean difference, -0.4%; $p < 0.01$). There were a large number of secondary end points. A portion of them were prespecified and analyses took into consideration the statistical impact of multiple comparisons; the remaining secondary outcomes were considered descriptive and p values were not reported. Among the prespecified secondary outcomes, treatment satisfaction (measured by the Diabetes Treatment Satisfaction Questionnaire [DTSQ]) was significantly higher in the CGM phase than in the conventional treatment phase ($p < 0.001$). Hypoglycemia outcomes were secondary descriptive outcomes. There was 1 (0.7%) severe hypoglycemic event during the CGM phase and 5 (3.5%) events during conventional therapy. The percentage of time with hypoglycemia (<70 mmol/L) was 2.8% during CGM treatment and 4.8% during conventional therapy.

In the second study, Beck et al (2017) randomized 158 patients on a 2:1 basis to 24 weeks of CGM ($n = 105$) or to usual care ($n = 53$). The trial included patients with type 1 diabetes who were ages 25 or older and had baseline HbA1c levels between 7.5% and 10%. Before randomization, patients underwent a 2-week period using a CGM system (without seeing data from the CGM) to ensure compliance. To be eligible, patients had to wear the CGM on at least 85% of days, calibrate the device at least twice daily and perform SMBG at least 3 times daily. The primary outcome (change in HbA1c levels at 24 weeks) was 1.0% in the CGM group and 0.4% in the

usual care group ($p < 0.001$), with a between-group difference of 0.6%. Prespecified secondary outcomes on the proportion of patients below a glycemic threshold at 24 weeks also favored the CGM group. The proportion of patients with HbA1c levels less than 7.0% was 18 (18%) in the CGM group and 2 (4%) in the control group ($p = 0.01$). The proportion of patients with HbA1c levels less than 7.5% was 39 (38%) in the CGM group and 6 (11%) in the control group ($p < 0.001$). Moreover, prespecified secondary outcomes related to hypoglycemia also differed significantly between groups, favoring the CGM group. The time spent in hypoglycemia less than 70 mg/dL was 43 minutes per day in the CGM group and 80 minutes per day in the usual care group ($p = 0.002$). Comparable numbers for time spent at less than 50 mg/dL were 6 minutes per day in the CGM group and 20 minutes per day in the usual care group ($p = 0.001$). The median change in the rate per 24 hours of hypoglycemia events lasting at least 20 minutes at less than 3.0 mmol/L (54 mg/dL) fell by 30% from 0.23 at baseline to 0.16 during follow-up in the CGM group but was practically unchanged (0.31 at baseline and 0.30 at follow-up) in the usual care group ($p = 0.03$). Quality of life measures assessing overall well-being (WHO-5), health status (EQ-5D-5L), diabetes distress (DDS), hypoglycemic fear (worry subscale of the HFS-II), and hypoglycemic confidence (HCS) have also been reported. There were no significant differences between CGM and usual care in changes in well-being, health status, or hypoglycemic fear. The CGM group demonstrated a greater increase in hypoglycemic confidence ($p = 0.01$) and a greater decrease in diabetes distress ($p = 0.01$) than the usual care group.

Pregnant Women

One trial of real-time CGM in pregnant women with type 1 diabetes has been reported. Study design results and gaps are summarized here and in Tables 3 to 6. Feig et al (2017) reported results of 2 multicenter RCTs in women ages 18 to 40 with type 1 diabetes who were receiving intensive insulin therapy and who were either pregnant (<13 weeks and 6 days of gestation) or planning a pregnancy.²⁰ The trial enrolling pregnant women is reviewed here. Women were eligible if they had a singleton pregnancy and HbA1c levels between 6.5% and 10.0%. The trial was conducted at 31 hospitals in North America and Europe. Women were randomized to CGM (Guardian REAL-Time or MiniMed Minilink system) plus capillary glucose monitoring or capillary glucose monitoring alone. Women in the CGM group were instructed to use the devices daily. Women in the control group continued their usual method of capillary glucose monitoring. The target glucose range was 3.5 to 7.8 mmol/L and target HbA1c levels were 6.5% or less in both groups. The primary outcome was the difference in change in HbA1c levels from randomization to 34 weeks of gestation. The proportion of completed scheduled study visits was high in both groups; however, participants using CGM had more unscheduled contacts, which were attributed both to sensor issues and to sensor-related diabetes management issues. The median frequency of CGM use was 6.1 days per week (interquartile range, 4.0-6.8) and 70% of pregnant participants used CGM for more than 75% of the time. The between-group difference in the change in HbA1c levels from baseline to 34 weeks of gestation was statistically significant favoring CGM (MD = -0.19%; 95% CI, -0.34 to -0.03; $p = 0.02$). Women in the CGM group spent an increased percentage of time in the recommended glucose control target range at 34 weeks of gestation (68% vs 61%, $p = 0.003$). There were no between-group differences in maternal hypoglycemia, gestational weight gain, or total daily insulin dose. A smaller proportion of infants of mothers in the CGM group were large-for-gestational age (odds ratio [OR], 0.51; 95% CI, 0.28 to 0.90; $p = 0.02$). In addition, for infants of mothers in the CGM group, there were fewer neonatal intensive care admissions lasting more than 24 hours (OR=0.48; 95% CI, 0.26 to 0.86;

p=0.02), fewer incidences of neonatal hypoglycemia requiring treatment with intravenous dextrose (OR=0.45, 0.22 to 0.89; p=0.025), and reduced total length of hospital stay (3.1 days vs 4.0 days; p=0.0091). Skin reactions occurred in 49 (48%) of 103 CGM participants and 8 (8%) of 104 control participants.

Table 3. RCT Characteristics for Real-Time CGM in Pregnant Women with Type 1 Diabetes

Study: Registration	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Feig et al (2017); NCT01788527	Canada, England, Scotland, Spain, Italy, Ireland, U.S.	31	2013-2016	Pregnant women (<14 wk gestation) with type 1 diabetes receiving intensive insulin therapy with HbA _{1c} levels between 6.5% and 10.0% (mean, 6.9%); mean age, 31 y	CGM (real-time, continuous) (n=108)	SMBG (n=107)

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c; RCT: randomized controlled trial; SMBG: self-monitored blood glucose.

Table 4. RCT Outcomes for Real-Time CGM in Pregnant Women with Type 1 Diabetes

Study	Infant			Maternal		
	Large-for-Gestational Age	Gestational Age at Delivery, wk	Severe Hypoglycemia	Caesarean Section	HbA _{1c} Levels: Change From Baseline to 34 Wk of Gestation	Severe Hypoglycemia
Feig et al (2017)						
n	211	201	200	202	173	214
CGM	53 (53%)	Median, 37.4	15 (15%)	63 (63%)	-0.54	11 (11%)
Control	69 (69%)	Median, 37.3	28 (28%)	74 (73%)	-0.35	12 (12%)
TE (95% CI)	OR=0.51 (0.28 to 0.90)	NR	OR=0.45 (0.22 to 0.89)	NR	-0.19% (-0.34% to -0.03%)	NR
p	0.02	0.50	0.025	0.18	0.02	1.0

Values are n or n (%) or as otherwise indicated.

CI: confidence interval; CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c; NR: not reported; OR: odds ratio; RCT: randomized controlled trial; TE: treatment effect.

Table 5. Relevance Gaps of RCTs for Real-Time CGM in Pregnant Women with Type 1 Diabetes

Study	Population	Intervention	Comparator	Outcomes	Follow-Up
Feig et al (2017)	4. Run-in period requirement may have biased selection to highly compliant participants	3. More unscheduled contacts in CGM group	3. More unscheduled contacts in CGM group	None noted	None noted
Key	1. <u>Intended use population unclear</u> 2. <u>Clinical context for treatment is unclear</u> 3. <u>Study population unclear</u> 4. <u>Study population not representative of intended</u>	1. <u>Not clearly defined</u> 2. <u>Version used unclear</u> 3. <u>Delivery not similar as comparator</u>	1. <u>Not clearly defined</u> 2. <u>Not standard or optimal</u> 3. <u>Delivery not similar intensity as intervention</u> 4. <u>Not delivered</u>	1. <u>Key health outcomes not addressed</u> 2. <u>Physiologic measures, not validated surrogates</u> 3. <u>Not CONSORT reporting of harms</u> 4. <u>Not established and validated</u>	1. <u>Not sufficient duration for benefits</u> 2. <u>Not sufficient duration for harms</u>

- use
- 5. Study population is subpopulation of intended use
- 4. Not delivered effectively
- effectively
- measurements
- 5. Clinically significant difference not prespecified
- 6. Clinically significant difference not supported

CGM: continuous glucose monitoring; RCT: randomized controlled trial.

Table 6. Study Design and Conduct Gaps of RCTs for Real-Time CGM in Pregnant Women with Type 1 Diabetes

<u>Study</u>	<u>Allocation</u>	<u>Blinding</u>	<u>Selective Reporting</u>	<u>Follow-Up</u>	<u>Power</u>	<u>Statistical</u>
<u>Feig et al (2017)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>3, 4. Treatment effects and confidence intervals not calculated for some outcomes</u>
Key	<ul style="list-style-type: none"> 1. <u>Participants not randomly allocated</u> 2. <u>Allocation not concealed</u> 3. <u>Allocation concealment unclear</u> 4. <u>Inadequate control for selection bias</u> 	<ul style="list-style-type: none"> 1. <u>Not blinded to treatment assignment</u> 2. <u>Not blinded outcome assessment</u> 3. <u>Outcome assessed by treating physician</u> 	<ul style="list-style-type: none"> 1. <u>Not registered</u> 2. <u>Evidence of selective reporting</u> 3. <u>Evidence of selective publication</u> 	<ul style="list-style-type: none"> 1. <u>High loss to follow-up or missing data</u> 2. <u>Inadequate handling of missing data</u> 3. <u>High number of crossovers</u> 4. <u>Inadequate handling of crossovers</u> 5. <u>Inappropriate exclusions</u> 6. <u>Not intent to treat analysis (per protocol for noninferiority trials)</u> 	<ul style="list-style-type: none"> 1. <u>Power calculations not reported</u> 2. <u>Power not calculated for primary outcome</u> 3. <u>Power not based on clinically important difference</u> 	<ul style="list-style-type: none"> 1. <u>Test is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event</u> 2. <u>Test is not appropriate for multiple observations per patient</u> 3. <u>Confidence intervals and/or p values not reported</u> 4. <u>Comparative treatment effects not calculated</u>

CGM: continuous glucose monitoring; RCT: randomized controlled trial.

Section Summary: CGM Devices for Long-Term Use in Type 1 Diabetes

Numerous RCTs and several systematic reviews of RCTs have evaluated CGM in patients with type 1 diabetes. A 2017 individual patient data analysis, using data from 11 RCTs, found that reduction in HbA1c levels was significantly greater with real-time CGM compared with a control intervention. In addition, a 2012 meta-analysis of 6 RCTs found a significantly larger decline in HbA1c levels 6 months in CGM users than the SMBG group. There are few studies beyond 6 months. Two recent RCTs in patients who used multiple daily insulin injections and were highly compliant with CGM devices during run-in phases found that CGM was associated with a larger reduction in HbA1c levels than previous studies. Reductions were 0.4% and 0.6%, respectively, compared with approximately 0.2 to 0.3% in previous analyses. One of the 2 RCTs prespecified hypoglycemia-related outcomes, and time spent in hypoglycemia was significantly less in the CGM group.

One RCT in pregnant women with type 1 diabetes (n=215) has compared CGM with SMBG. Adherence was high in the CGM group. The difference in the change in HbA1c levels from baseline to 34 weeks of gestation was statistically significant favoring CGM, and women in the CGM group spent an increased percentage of time in the recommended glucose control target range at 34 weeks of gestation. There were no between-group differences in maternal hypoglycemia, gestational weight gain, or total daily insulin dose. A smaller proportion of infants of mothers in the CGM group were large for gestational age, had neonatal intensive care admissions lasting more than 24 hours, and had neonatal hypoglycemia requiring treatment. The total length of hospital stay was shorter by almost 1 day in the CGM group.

CGM Devices for Intermittent, Short Term use

The meta-analyses of CGM devices for type 1 diabetes tended to combine studies of intermittent glucose monitoring with studies of continuous long term monitoring. For this body of evidence, there is variability in the definitions of intermittent monitoring and the specific monitoring protocols used. In addition, many of the trials of intermittent monitoring included additional interventions to optimize glucose control such as education and recommendations on lifestyle modifications.

Two meta-analyses were identified that reported separate subgroup analysis for intermittent monitoring. In the 2012 Cochrane review, there were 4 studies (216 patients) comparing real-time intermittent CGM systems to SMBG, and the pooled effect estimate for change in HbA1c at 3 months was not statistically significant (MD change = -0.18; 95% CI, -0.42 to 0.05). The 2011 meta-analysis of RCTs on CGM described previously also included a separate analysis of 8 RCTs of intermittent monitoring. On pooled analysis, there was a statistically significant reduction in HbA1c with CGM compared with SMBG (WMD = -0.26; 95% CI, -0.45 to -0.06).

The largest individual RCT was the MITRE trial, published by Newman and colleagues in 2009, was conducted to evaluate whether the additional information provided by use of minimally invasive glucose monitors resulted in improved glucose control in patients with poorly controlled insulin-requiring diabetes. This was a 4-arm RCT conducted at secondary care diabetes clinics in 4 hospitals in England. In this study, 404 people aged older than 18 years, with insulin-treated diabetes mellitus (Types 1 or 2) for at least 6 months, who were receiving 2 or more injections of insulin daily, were eligible. The majority of participants, 57%, had Type 1 diabetes, 41% had

Type 2 diabetes and 2% were classified as “other.” Participants had two HbA1c values of at least 7.5% in the 15 months prior to entry and were randomized to 1 of 4 groups. Two groups received minimally invasive glucose monitoring devices (GlucoWatch Biographer or MiniMed Continuous Glucose Monitoring System, CGMS). Intermittent CGM was used (i.e., monitoring was performed over several days at various points in the study). These groups were compared with an attention control group (standard treatment with nurse feedback sessions at the same frequency as those in the device groups) and a standard control group (reflecting common practice in the clinical management of diabetes). Change in HbA1c from baseline to 3, 6, 12, and 18 months was the primary indicator of short- to long-term efficacy in this study. At 18 months, all groups demonstrated a decline in HbA1c levels from baseline. Mean percentage changes in HbA1c were -1.4 for the GlucoWatch group, -4.2 for the CGMS group, -5.1 for the attention control group, and -4.9 for the standard care control group. In the intent-to-treat (ITT) analysis, no significant differences were found between any of the groups at any of the assessment times. There was no evidence that the additional information provided by the devices resulted in any change in the number or nature of treatment recommendations offered by the nurses. Use and acceptability indicated a decline in use of both devices, which was most marked in the GlucoWatch group by 18 months (20% still using GlucoWatch vs. 57% still using the CGMS). In this study of unselected patients, use of continuous glucose monitors (CGMS on an intermittent basis) did not lead to improved clinical outcomes.

Pregnant Women

Systematic Reviews

Voormolen et al (2013) published a systematic review of the literature on CGM during pregnancy. They identified 11 relevant studies (total N=534 women). Two were RCTs, one of which was the largest of the studies (N=154). Seven studies used CGMs that do not have data available in real-time; the remaining 4 studies used real-time CGM. Reviewers did not pool study findings; they concluded that the evidence was limited on the efficacy of CGM during pregnancy. The published RCTs are described next.

Randomized Controlled Trials

Two RCTs of intermittent glucose monitoring in pregnant women with type 1 or type 2 diabetes are summarized in Tables 7 to 10 and the following paragraphs. While both trials included a mix of women with type 1 and type 2 diabetes, most women had type 1 diabetes in both trials, so the trials are reviewed in this section.

In 2008, Murphy et al in the U.K. randomized 71 pregnant women with Type 1 (n=46) or Type 2 (n=25) diabetes to CGM or usual care. The intervention consisted of up to 7 days of CGM at intervals of 4 to 6 weeks between 8 and 32 weeks' gestation. Neither participants nor physicians had access to the measurements during sensor use; data were reviewed at study visits. In addition to CGM, the women were advised to measure blood glucose levels at least seven times a day. Baseline HbA1c was 7.2% (standard deviation [SD] =0.9) in the CGM group and 7.4% (SD=1.5) in the usual care group. The primary study outcome was maternal glycemic control during the second and third trimesters. Eighty percent of women in the CGM group wore the monitor at least once per trimester. Mean HbA1c levels were consistently lower in the intervention arm, but differences between groups were statistically significant only at week 36. For example, between 28 and 32 weeks' gestation, mean HbA1c levels were 6.1% (SD=0.60 in the CGM group and

6.4% (SD=0.8) in the usual care group, p=0.10. The prevalence of large-for-gestational age infants (at least 90th percentile) was a secondary outcome. Thirteen of 37 (35%) infants in the CGM group were large-for-gestational age compared with 18 of 30 (60%) in the usual care group. The odds ratio for reduced risk of a large-for-gestational age infant with CGM was 0.36 (95% CI, 0.13 to 0.98, p=0.05).

Secher et al (2013) randomized 154 women with type 1 (n=123) and type 2 (n=31) diabetes to real-time CGM in addition to routine pregnancy care (n=79) or routine pregnancy care alone (n=75). Patients in the CGM group were instructed to use the CGM device for 6 days before each of 5 study visits and were encouraged to use the devices continuously; 64% of participants used the devices per-protocol. Participants in both groups were instructed to perform 8 daily self-monitored plasma glucose measurements for 6 days before each visit. Baseline mean HbA1c levels were 6.6% in the CGM group and 6.8% in the routine care group. The 154 pregnancies resulted in 149 live births and 5 miscarriages. The prevalence of large-for-gestational age infants (at least 90th percentile), the primary study outcome, was 45% in the CGM group and 34% in the routine care group. The difference between groups was not statistically significant (p=0.19). Also, no statistically significant differences were found between groups for secondary outcomes, including the prevalence of preterm delivery and the prevalence of severe neonatal hypoglycemia. Women in this trial had low baseline HbA1c levels, which might explain the lack of impact of CGM on outcomes. Other factors potentially contributing to the negative findings included the intensive SMBG routine in both groups and the relatively low compliance rate in the CGM group.

Table 7. Key RCT Characteristics for Intermittent CGM in Pregnant Women with Type 1 Diabetes

<u>Author:</u> <u>Registration</u>	<u>Countries</u>	<u>Sites</u>	<u>Dates</u>	<u>Participants</u>	<u>Interventions</u>	
					<u>Active</u>	<u>Comparative</u>
Murphy et al (2008); ISRCTN844615 81	U.K.	2	2003- 2006	Pregnant women with type 1 (65%) and type 2 (35%) diabetes; mean gestational age, 9.2 wk; mean HbA _{1c} level, 7.3%; mean age, 31 y	CGM (up to 7 d of CGM at intervals of 4-6 wk) plus SOC (n=38)	SOC (n=33)
Secher et al (2013); NCT00994357	Denmark	1	2009- 2011	Pregnant women with type 1 (80%) or type 2 (20%) diabetes; mean gestational age, <14 wk; median HbA _{1c} level, 6.7%; median age, 32 y	CGM (for 6 d before each study visits; encouraged to used continuously) plus routine care (n=79)	Routine care (n=75)

CGM: continuous glucose monitoring; HbA1c: hemoglobin A1c; RCT: randomized controlled trial; SOC: standard of care.

Table 8. RCT Outcomes of Intermittent CGM in Pregnant Women with Type 1 Diabetes

Study	Large-for-Gestational Age	Infant			Maternal	
		Gestational Age at Delivery Weeks	Severe Hypoglycemia ^a	Caesarean Section	HbA _{1c} Levels: At 36 Wk ^a Gestation ^a	Severe Hypoglycemia ^a
<u>Murphy et al (2008)</u>						
<u>n</u>	<u>71</u>	<u>71</u>	<u>68</u>	<u>69</u>	<u>71</u>	<u>NR</u>
<u>CGM</u>	<u>13 (35%)</u>	<u>Mean, 37.6</u>	<u>3 (8%)</u>	<u>27 (71%)</u>	<u>Mean, 5.8%</u>	
<u>Control</u>	<u>18 (60%)</u>	<u>Mean, 37.5</u>	<u>5 (17%)</u>	<u>21 (61%)</u>	<u>Mean, 6.4%</u>	
<u>TE (95% CI)</u>	<u>OR=0.36 (0.13 to 0.98)</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>0.6% (CI NR)</u>	
<u>p</u>	<u>0.05</u>	<u>0.80</u>	<u>0.50</u>	<u>0.40</u>	<u>0.007</u>	
Days						
<u>Secher et al (2013)</u>						
<u>n</u>	<u>154</u>	<u>154</u>	<u>145</u>	<u>154</u>		<u>154</u>
<u>CGM</u>	<u>34 (45%)</u>	<u>Median, 263</u>	<u>9 (13%)</u>	<u>28 (37%)</u>	<u>Median, 6.0%</u>	<u>16%</u>
<u>Control</u>	<u>25 (34%)</u>	<u>Median, 264</u>	<u>10 (14%)</u>	<u>33 (45%)</u>	<u>Median, 6.1%</u>	<u>16%</u>
<u>TE (95% CI)</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>p</u>	<u>0.19</u>	<u>0.14</u>	<u>0.88</u>	<u>0.30</u>	<u>0.63</u>	<u>0.91</u>

Values are n or n (%) or as otherwise indicated.

CGM: continuous glucose monitoring; CI: confidence interval; HbA_{1c}: hemoglobin A_{1c}; NR: not reported; OR: odds ratio;

RCT: randomized controlled trial; TE: treatment effect.

^a N inconsistently reported for HbA_{1c} outcome.

In summary, 2 studies of intermittent glucose monitoring conducted in Europe included pregnant women with type 1 or 2 diabetes, with most having type 1 diabetes. Murphy et al (2008) included intermittent, retrospective monitoring with CGM; Secher et al (2013) included intermittent, real-time monitoring. The intervention started in early pregnancy in these studies; mean age was in the early thirties and mean baseline HbA_{1c} level was greater than 6.5%. There was no statistically significant difference between CGM and routine care for maternal HbA_{1c} levels at 36 weeks in Secher; the difference in HbA_{1c} levels at 36 weeks was about 0.6% (p=0.007) in Murphy. Secher also reported no difference in severe maternal hypoglycemia. The proportion of infants that were large for gestational age (>90th percentile) was higher in the CGM group in Secher, although not statistically significantly higher; the difference in large for gestational age was statistically significantly lower for CGM in Murphy. The differences in the proportions of infants born via caesarean section, gestational age at delivery, and infants with severe hypoglycemia were not statistically significant in either trial.

Table 9. Relevance Gaps of RCTs of Intermittent CGM in Pregnant Women with Type 1 Diabetes

<u>Study</u>	<u>Population</u>	<u>Intervention</u>	<u>Comparator</u>	<u>Outcomes</u>	<u>Follow-Up</u>
<u>Murphy et al (2008)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>
<u>Secher et al (2013)</u>	<u>4. Study population had relatively low HbA_{1c}</u>	<u>4. Only 64% of the participants used devices per protocol</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>
<u>Key</u>	<u>6. Intended use population unclear</u> <u>7. Clinical context for treatment is unclear</u> <u>8. Study population unclear</u> <u>9. Study population not representative of intended use</u> <u>10. Study population is subpopulation of intended use</u>	<u>5. Not clearly defined</u> <u>6. Version used unclear</u> <u>7. Delivery not similar intensity as comparator</u> <u>8. Not delivered effectively</u>	<u>5. Not clearly defined</u> <u>6. Not standard or optimal</u> <u>7. Delivery not similar intensity as intervention</u> <u>8. Not delivered effectively</u>	<u>7. Key health outcomes not addressed</u> <u>8. Physiologic measures, not validated surrogates</u> <u>9. Not CONSORT reporting of harms</u> <u>10. Not established and validated measurements</u> <u>11. Clinically significant difference not prespecified</u> <u>12. Clinically significant difference not supported</u>	<u>3. Not sufficient duration for benefits</u> <u>4. Not sufficient duration for harms</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c; RCT: randomized controlled trial.

Table 10. Study Design and Conduct Gaps of RCTs of Intermittent Glucose Monitoring in Pregnant Women with Type 1 Diabetes

<u>Study</u>	<u>Allocation</u>	<u>Blinding</u>	<u>Selective Reporting</u>	<u>Follow-Up</u>	<u>Power</u>	<u>Statistical</u>
<u>Murphy et al (2008)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>3, 4. Treatment effects and confidence intervals not calculated for some outcomes</u>
<u>Secher et al (2013)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>3, 4. Treatment effects and confidence intervals not calculated</u>
<u>Key</u>	<u>1. Participants not randomly allocated</u> <u>2. Allocation not concealed</u> <u>3. Allocation concealment</u>	<u>1. Not blinded to treatment assignment</u> <u>2. Not blinded outcome assessment</u> <u>3. Outcome</u>	<u>1. Not registered</u> <u>2. Evidence of selective reporting</u> <u>3. Evidence of selective publication</u>	<u>1. High loss to follow-up or missing data</u> <u>2. Inadequate handling of missing data</u> <u>3. High number of crossovers</u> <u>4. Inadequate handling of</u>	<u>1. Power calculations not reported</u> <u>2. Power not calculated for primary outcome</u> <u>3. Power not based</u>	<u>1. Test is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event</u> <u>2. Test is not appropriate for multiple observations per patient</u> <u>3. Confidence intervals and/or</u>

unclear
4. Inadequate
control for
selection bias

assessed by
treating
physician

crossovers
5. Inappropriate
exclusions
6. Not intent to treat
analysis (per protocol
for noninferiority trials)

on clinically
important
difference

p-values not reported
4. Comparative treatment
effects not calculated

CGM: continuous glucose monitoring; RCT: randomized controlled trial.

Section Summary: Glucose Monitoring Devices for Intermittent, Short-Term Use in Type 1 Diabetes

For intermittent, short-term monitoring of type 1 diabetes, there are few RCTs and systematic reviews. Some trials have reported improvements in glucose control for the intermittent monitoring group, but limitations in this body of evidence preclude conclusions. The definitions of intermittent control and the specific monitoring protocols varied. In some studies, intermittent monitoring was part of a larger strategy aimed at optimizing glucose control, and the impact of monitoring cannot be separated from the impact of other interventions.

Two RCTs of intermittent glucose monitoring have been conducted in pregnant women with both type 1 and 2 diabetes, with most having type 1 diabetes. One study reported a difference in HbA1c levels at 36 weeks and the proportion of infants that were large for gestational age (>90th percentile) favoring CGM while the second study did not. The differences in the proportions of infants born via caesarean section, gestational age at delivery, and infants with severe hypoglycemia were not statistically significant in either study.

Type 2 Diabetes for Long and Short term Glucose Monitoring

The analysis of type 2 diabetes does not distinguish between indications 3 and 4 (long-term and short-term glucose monitoring), consistent with the literature.

Systematic Reviews

Two systematic reviews (previously described) also reported on the efficacy of CGM in patients with type 2 diabetes. A comparison of the trials of type 2 diabetes included in the systematic reviews and meta-analyses in these reviews is shown in Table 11.

Table 11. Comparison of CGM Trials for Type 2 Diabetes Included in Systematic Reviews

<u>Primary Study</u>	<u>Gandhi et al (2011)</u>	<u>Poolsup et al (2013)</u>
<u>Allen et al (2008)</u>	●	●
<u>Yoo et al (2008)</u>	●	●
<u>Cosson et al (2009)</u>	●	●
<u>Ehrhardt et al (2011)</u>		●

CGM: continuous glucose monitoring

A summary of the characteristics of the systematic reviews is shown in Table 12. Results are briefly described in Table 13 and the following. Gandhi et al (2011) discussed above, included studies of Type 1 and Type 2 diabetes. Three RCTs included patients with Type 2 diabetes. One RCT included patients with either type of diabetes. There was a mixture of patients with Type 2 diabetes who did and did not require insulin. Two of the 3 trials evaluated retrospective intermittent CGM of different lengths and durations and the third evaluated real-time intermittent CGM. Patients included in the studies had baseline HbA1c > 8%. In a meta-analysis of the three trials, there was a statistically significant reduction in HbA1C with CGM compared to SMBG in adults with Type 2 diabetes (WMD: -0.70, 95% CI: -1.14 to -0.27). In 2013, Poolsup et al conducted a meta-analysis of four trials using adults with Type 2 diabetes. Three of the trials in Poolsup overlapped with Gandhi; the remaining trial also evaluated real-time CGM but with a longer period of use (2 weeks on and 1 week off for 3 months). In a pooled analysis, CGM had greater efficacy in terms of HbA1c than usual care. The pooled mean difference in HbA1c was -

0.31% 95% CI, -0.6 to 0.02, p=0.04). Because of a lack of statistical heterogeneity among studies, subgroup analyses (e.g., by type of CGM device) were not performed.

Table 12. Systematic Review Characteristics for CGM in Type 2 Diabetes

<u>Study</u>	<u>Dates</u>	<u>Trials</u>	<u>Participants</u>	<u>N (Range)</u>	<u>Design</u>	<u>Duration</u>
<u>Gandhi et al (2011)</u>	<u>1996-2010</u>	<u>3</u>	<u>Adult outpatients with T2D; mean baseline HbA_{1c} level >8%</u>	<u>128 (25-57)</u>	<u>RCT</u>	<u>At least 8 wk (median, 3 mo.)</u>
<u>Poolsup et al (2013)</u>	<u>Up to 2013</u>	<u>4</u>	<u>Adults with T2D</u>	<u>228 (25-100)</u>	<u>RCT</u>	<u>At least 8 wk (median, 3 mo.)</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c; RCT: randomized controlled trial; T2D: type 2 diabetes.

Table 13. Meta-Analytic Results for CGM in Type 2 Diabetes

<u>Study</u>	<u>Reduction in HbA_{1c} Levels (Mean Difference)</u>	<u>Hypoglycemic Events</u>	<u>Diabetes Complications (retinopathy, nephropathy, neuropathy, diabetic foot)</u>	<u>Health-Related Quality of Life</u>
<u>Gandhi et al (2011)</u>				
<u>Total N</u>	<u>128</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>PE (95% CI)</u>	<u>-0.70 (-1.14 to -0.27)</u>			
<u>p</u>	<u>NR</u>			
<u>I²</u>	<u>0%</u>			
<u>Poolsup et al (2013)</u>				
<u>Total N</u>	<u>228</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>PE (95% CI)</u>	<u>-0.31 (-0.60 to -0.02)</u>			
<u>p</u>	<u>0.04</u>			
<u>I²</u>	<u>0%</u>			

CGM: continuous glucose monitoring; CI: confidence interval; HbA_{1c}: hemoglobin A1c; NR: not reported; PE: pooled effect.

Randomized Controlled Trials

Several RCTs of CGM in adults with type 2 diabetes are summarized in Tables 14 to 17. The largest and most recent studies are also briefly summarized in the following paragraphs. The studies were conducted in North America, Europe, and Asia. Baseline HbA_{1c} levels were between 8.5% and 9.0% in the RCTs, with participants having a mean baseline age range in the mid-50s and early-60s. The RCTs used a mixed of intermittent and continuous, real-time monitoring.

Ehrhardt and colleagues published 2 reports (2011, 2012) from an RCT evaluating the largest sample (N=100) in the Poolsup et al (2013) systematic review (accounting for 45% of the weight in the pooled analysis of HbA_{1c} levels). The trial evaluated intermittent use of a CGM device in adult patients with Type 2 diabetes who were treated with diet/exercise and/or glycemic lowering medications but not prandial insulin and had an initial HbA_{1c} of at least 7% but not more than 12%. The study compared real-time continuous monitoring with the DexCom device used for four 2-week cycles (2 weeks on/ 1 week off) to self-monitoring of blood glucose (SMBG). The primary efficacy outcome was mean change in HbA_{1c}. Mean (SD) HbA_{1c} in the CGM group were 8.4% (1.5%) at baseline, 7.4% (1.0%) at 12 weeks, 7.3% (1.1%) at 24 weeks, and 7.7% (1.1%) at 52 weeks. In the SMBG group, these values (SD) were 8.2% (1.1%) at baseline, 7.7% (1.2%) at 12 weeks, 7.6% (1.3%) at 24 weeks, and 7.9% (1.4%) at 52 weeks. Over the course of the study, the reduction in HbA_{1c} was significantly greater than in the SMBG group (p=0.04).

After adjusting for potential confounding variables including age, sex, baseline therapy, and whether the individual started taking insulin during the study, the difference between groups over time remained statistically significant ($p < 0.0001$). The investigators also evaluated SMBG results from both groups. The mean proportion of SMBG tests that were less than 70 mg/dL were 3.6% in the CGM group and 2.5% in the SMBG group ($p = 0.06$).

The RCT by Sato et al (2016), included 34 patients with type 2 diabetes who were at least 20 years old and were on insulin injection therapy, had HbA1c levels between 6.9% and 11.0% during the previous 3 months, with fluctuations of HbA1c within 0.5%. All patients conducted SMBG and used a retrospective CGM device for 4 to 5 days before each of 3 clinic visits, 2 months apart. At the clinic visits, patients' were evaluated and suggestions were made to improve glucose control by lifestyle changes and secondarily by changing medication doses. In the intervention group, but not the control group, the patients and physicians had access to the CGM data at the clinic visits. The primary endpoint was change in HbA1c and this did not differ significantly between groups at the end of the study, compared with baseline, between the first and second visit or between the second and third visits. HbA1c changed little in either group. In the intervention group, the mean (SD) baseline HbA1c was 8.2% (1.2%) and the mean final HbA1c was also 8.2% (SD: 1.3%). Comparable percentages in the control group were 8.2% (0.9%), and 7.9% (0.8%). In this study, which was conducted in Japan, decisions around medication doses were made only by the physician at clinic visits and practices may differ in other countries.

The largest RCT, Multiple Daily Injections and Continuous Glucose Monitoring in Diabetes (DIAMOND), was reported by Beck et al (2017). DIAMOND was performed at 25 endocrinology practices in North America (22 in the United States, 3 in Canada) and enrolled adults with type 2 diabetes receiving multiple daily injections of insulin. One-hundred fifty-eight patients were randomized into 2 groups, CGM and usual care (n=79 in each group). Patients compliant during a run-in period were eligible for randomization. Patients in both groups were given a blood glucose meter. Participants in the CGM group were given a Dexcom G4 Platinum CGM System (Dexcom) and instructions on use. Change in HbA1c level from baseline to 24 weeks was the primary outcome. Analyses were adjusted for baseline HbA1c levels and clinic were performed using intention-to-treat analysis with missing data handling by multiple imputation. At baseline, the mean total daily insulin dose was 1.1 U/kg/d. Week 24 follow-up was completed by 97% of the CGM group and 95% of the control group. Mean CGM use was greater than 6 d/wk at 1 month, 3 months, and 6 months. The adjusted difference in mean change in HbA1c level from baseline to 24 weeks was -0.3% (95% CI, -0.5% to 0.0%; $p = 0.022$) favoring CGM. The adjusted difference in the proportion of patients with a relative reduction in HbA1c level of 10% or more was 22% (95% CI, 0% to 42%; $p = 0.028$) favoring CGM. There were no events of severe hypoglycemia or diabetic ketoacidosis in either group. The treatment groups did not differ in any of the quality of life measures.

Table 14. RCT Characteristics for Glucose Monitoring in Type 2 Diabetes

<u>Study:</u> <u>Registration</u>	<u>Countries</u>	<u>Sites</u>	<u>Dates</u>	<u>Participants</u>	<u>Interventions</u>	
					<u>Active</u>	<u>Comparat or</u>
<u>Allen et al (2008)</u>	<u>U.S.</u>	<u>2</u>	<u>NR</u>	<u>Adults with T2D not receiving insulin with HbA_{1c} levels >7.5% (baseline mean, 8.6%), not participating in physical activity; mean age, 57 y</u>	<u>Diabetes education plus CGM for 3 d (n=27)</u>	<u>Diabetes education (n=25)</u>
<u>Yoo et al (2008)</u>	<u>Korea</u>	<u>4</u>	<u>2007</u>	<u>Adults with T2D using oral antidiabetic agents or insulin with HbA_{1c} levels 8.0%-10.0% (baseline mean, 9%); mean age, 56 y</u>	<u>CGM (3 d at a time for 3 mo.) (n=32)</u>	<u>SMBG (n=33)</u>
<u>Cosson et al (2009)</u>	<u>France</u>	<u>5</u>	<u>NR</u>	<u>Adults with T1D or T2D treated with oral antidiabetic agents with or without insulin with HbA_{1c} levels 8.0%-10.5% (baseline mean, 9.1% in T2D); mean age, 57 y in T2D</u>	<u>CGM for 48 h at baseline and 3 mo.; CGM data shared with physician and patient (n=11 with T2D)</u>	<u>“Blinded” CGM (n=14 in T2D)</u>
<u>Ehrhardt et al (2011)</u>	<u>U.S.</u>	<u>1</u>	<u>NR</u>	<u>Adults with T2D using oral antidiabetic agents without prandial insulin with HbA_{1c} levels 7.0%-12.0% (baseline mean, 8.3%), mean age, 58 y</u>	<u>Real-time CGM for 4 cycles of 3 wk (n=50)</u>	<u>SMBG (n=50)</u>
<u>Sato et al (2016); UMIN: 000012034^a</u>	<u>Japan</u>	<u>1</u>	<u>2012-2014</u>	<u>Adults with T2D using insulin with HbA_{1c} levels 6.9%-11.0% (baseline mean, 8.2%); mean age, 62 y</u>	<u>CGM for 4-5 d every 4 mo.; reviewed at study visits (n=17)</u>	<u>“Blinded” CGM (n=17)</u>
<u>Beck et al (2017) (DIAMOND); NCT02282397</u>	<u>U.S., Canada</u>	<u>25</u>	<u>2014-2016</u>	<u>Adults with T2D using multiple daily injections of insulin with HbA_{1c} levels 7.5%-10.0% (baseline mean, 8.5%); mean age, 60 y</u>	<u>Real-time CGM, continuous, real-time (n=79)</u>	<u>SMBG (n=79)</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A_{1c}; NR: not reported; SMBG: self-monitored blood glucose; T1D: type 1 diabetes; T2D: type 2 diabetes.

^a Registered with the University Hospital Medical Information Network in Japan.

Most RCTs used a type of intermittent monitoring; some reported data for patients in real-time while others provided data reviewed only at study visits. Four of the 6 RCTs of CGM in type 2 diabetes reported a statistically significant larger decrease in HbA_{1c} levels with CGM than with control. In Cosson et al (2009), the comparative treatment effect was not reported, but the CGM group had a statistically significant reduction in HbA_{1c} levels from baseline to 3 months. Few other outcomes were reported. Beck et al (2017) reported more patients in CGM with a relative reduction in HbA_{1c} levels of greater than 10% at 24 weeks but no difference in the quality of life measures. No trials reported on follow-up beyond 6 months. Thus the effect of CGM on outcomes related to diabetic complications is unknown. Only 2 RCTs used blinded CGM; in one, there was no difference in reduction in HbA_{1c} levels between CGM and control.

Table 15. RCT Outcomes for Glucose Monitoring in Type 2 Diabetes

<u>Study</u>	<u>Reduction in HbA_{1c} Levels (Mean Range)</u>	<u>HbA_{1c} Level <7.0%, %</u>	<u>Relative Reduction in HbA_{1c} Level ≥10%, %</u>	<u>Hypoglycemic or Ketoacidosis Events</u>	<u>Diabetes Complications (retinopathy, nephropathy, neuropathy, diabetic foot)</u>	<u>Health-Related Quality of Life</u>
<u>Change From Baseline to 8 Wk</u>						
<u>Allen et al (2008)</u>						
<u>N</u>	<u>46</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>CGM</u>	<u>8.9% to 7.7%</u>					
<u>Control</u>	<u>8.4% to 8.1%</u>					
<u>TE (95% CI)</u>	<u>NR</u>					
<u>p</u>	<u><0.05</u>					
<u>Change From Baseline to 3 Mo</u>						
<u>Yoo et al (2008)</u>						
<u>N</u>	<u>57</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>CGM</u>	<u>9.1% to 8.0%</u>					
<u>Control</u>	<u>8.7% to 8.3%</u>					
<u>TE (95% CI)</u>	<u>NR</u>					
<u>p</u>	<u>0.004</u>					
<u>Time Spent with Hypoglycemia, min</u>						
<u>Cosson et al (2009)</u>						
<u>N</u>	<u>25</u>	<u>NR</u>	<u>NR</u>	<u>19</u>	<u>NR</u>	<u>NR</u>
<u>CGM</u>	<u>9.2% to 8.6%</u>			<u>18</u>		
<u>Control</u>	<u>9.0% to 8.8%</u>			<u>11</u>		
<u>TE (95% CI)</u>	<u>NR</u>			<u>NR</u>		
<u>Change From Baseline to 12 Wk</u>						
<u>Ehrhardt et al (2011)</u>						
<u>N</u>	<u>100</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>CGM</u>	<u>8.4% to 7.4%</u>					
<u>Control</u>	<u>8.2% to 7.7%</u>					
<u>TE (95% CI)</u>	<u>NR</u>					
<u>p</u>	<u>0.006</u>					

Study	Reduction in HbA _{1c} Levels (Mean Range)	HbA _{1c} Level <7.0%, %	Relative Reduction in HbA _{1c} Level ≥10%, %	Hypoglycemic or Ketoacidosis Events	Diabetes Complications (retinopathy, nephropathy, neuropathy, diabetic foot)	Health-Related Quality of Life
Sato et al (2016)						
<u>N</u>	<u>34</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>
<u>CGM</u>	<u>8.2% to 8.2%</u>					
<u>Control</u>	<u>8.2% to 7.9%</u>					
<u>TE (95% CI)</u>	<u>NR</u>					
<u>p</u>	<u>≥0.05</u>					
	<u>Change From Baseline to 24 Wk</u>	<u>At 24 Wk</u>	<u>At 24 Wk</u>			<u>DDS Overall Mean Score at 24 Wk</u>
Beck et al (2017)						
<u>N</u>	<u>158</u>	<u>158</u>	<u>158</u>	<u>158</u>	<u>NR</u>	<u>150</u>
<u>CGM</u>	<u>8.6% to 7.7%</u>	<u>11 (14%)</u>	<u>40 (52%)</u>	<u>0</u>		<u>1.8</u>
<u>Control</u>	<u>8.6% to 8.2%</u>	<u>9 (12%)</u>	<u>24 (32%)</u>	<u>0</u>		<u>1.8</u>
<u>TE (95% CI)</u>	<u>-0.3 (-0.5 to 0.0)</u>	<u>3% (-9% to 14%)</u>	<u>22% (0% to 42%)</u>			<u>NR</u>
<u>p</u>	<u>0.022</u>	<u>0.88</u>	<u>0.028</u>			

Table 16. Relevance Gaps of RCTs for Glucose Monitoring in Type 2 Diabetes

Study	Population	Intervention	Comparator	Outcomes	Follow-Up
<u>Allen et al (2008)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Focused on HbA_{1c}; did not include outcomes on adverse events, quality of life, or diabetic complications</u>	<u>1. Follow-up not sufficient to determine effects on diabetic complications</u>
<u>Yoo et al (2008)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Focused on HbA_{1c}; did not include outcomes on adverse events, quality of life, or diabetic complications</u>	<u>1. Follow-up not sufficient to determine effects on diabetic complications</u>
<u>Cosson et al (2009)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Focused on HbA_{1c}; did not include outcomes on adverse events, quality of life, or diabetic complications</u>	<u>1. Follow-up not sufficient to determine effects on diabetic complications</u>
<u>Ehrhardt et al</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Focused on HbA_{1c}; did not</u>	<u>1. Follow-up not sufficient to</u>

(2011)

include outcomes on adverse events, quality of life, or diabetic complications
6. No justification for clinically significant difference

determine effects on diabetic complications; patients reportedly followed for 52 wk but data not reported.

<u>Sato et al (2016)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Focused on HbA_{1c}; did not include outcomes on adverse events, quality of life, or diabetic complications</u>	<u>1. Follow-up not sufficient to determine effects on diabetic complications</u>
<u>Beck et al (2017) (DIAMOND)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Did not include outcomes on diabetic complications</u>	<u>1. Follow-up not sufficient to determine effects on diabetic complications</u>
Key	<ul style="list-style-type: none"> 1. <u>Intended use population unclear</u> 2. <u>Clinical context for treatment is unclear</u> 3. <u>Study population unclear</u> 4. <u>Study population not representative of intended use</u> 5. <u>Study population is subpopulation of intended use</u> 	<ul style="list-style-type: none"> 1. <u>Not clearly defined</u> 2. <u>Version used unclear</u> 3. <u>Delivery not similar intensity as comparator</u> 4. <u>Not delivered effectively</u> 	<ul style="list-style-type: none"> 1. <u>Not clearly defined</u> 2. <u>Not standard or optimal</u> 3. <u>Delivery not similar intensity as intervention</u> 4. <u>Not delivered effectively</u> 	<ul style="list-style-type: none"> 1. <u>Key health outcomes not addressed</u> 2. <u>Physiologic measures, not validated surrogates</u> 3. <u>Not CONSORT reporting of harms</u> 4. <u>Not established and validated measurements</u> 5. <u>Clinically significant difference not prespecified</u> 6. <u>Clinically significant difference not supported</u> 	<ul style="list-style-type: none"> 1. <u>Not sufficient duration for benefits</u> 2. <u>Not sufficient duration for harms</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A_{1c}; RCT: randomized controlled trial.

Table 17. Study Design and Conduct Gaps of RCTs for Glucose Monitoring in Type 2 Diabetes

<u>Study</u>	<u>Allocation</u>	<u>Blinding</u>	<u>Selective Reporting</u>	<u>Follow-Up</u>	<u>Power</u>	<u>Statistical</u>
<u>Allen et al (2008)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>1. Registration not reported</u>	<u>None noted</u>	<u>2, 3. Power not calculated a priori; convenience sample size</u>	<u>3, 4. Treatment effects and confidence intervals not calculated</u>
<u>Yoo et al (2008)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>1. Registration not reported</u>	<u>None noted</u>	<u>None noted</u>	<u>3, 4. Treatment effects and confidence intervals not calculated</u>
<u>Cosson et al</u>	<u>None noted</u>	<u>None noted</u>	<u>1. Registration</u>	<u>2. Unclear how missing data</u>	<u>1, 2, 3. No power</u>	<u>3, 4. Treatment effects and</u>

<u>(2009)</u>			<u>not reported</u>	<u>were handled in analyses</u>	<u>calculations</u>	<u>confidence intervals not calculated</u>
<u>Ehrhardt et al (2011)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>1. Registration not reported</u>	<u>None noted</u>	<u>3. No justification for difference used for power calculation</u>	<u>3, 4. Treatment effects and CIs not calculated</u>
<u>Sato et al (2016)</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>3, 4. Treatment effects and confidence intervals not calculated</u>
<u>Beck et al (2017) (DIAMOND)</u>	<u>None noted</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>
<u>Key</u>	<ul style="list-style-type: none"> 1. <u>Participants not randomly allocated</u> 2. <u>Allocation not concealed</u> 3. <u>Allocation concealment unclear</u> 4. <u>Inadequate control for selection bias</u> 	<ul style="list-style-type: none"> 1. <u>Not blinded to treatment assignment</u> 2. <u>Not blinded outcome assessment</u> 3. <u>Outcome assessed by treating physician</u> 	<ul style="list-style-type: none"> 1. <u>Not registered</u> 2. <u>Evidence of selective reporting</u> 3. <u>Evidence of selective publication</u> 	<ul style="list-style-type: none"> 1. <u>High loss to follow-up or missing data</u> 2. <u>Inadequate handling of missing data</u> 3. <u>High number of crossovers</u> 4. <u>Inadequate handling of crossovers</u> 5. <u>Inappropriate exclusions</u> 6. <u>Not intent to treat analysis (per protocol for noninferiority trials)</u> 	<ul style="list-style-type: none"> 1. <u>Power calculations not reported</u> 2. <u>Power not calculated for primary outcome</u> 3. <u>Power not based on clinically important difference</u> 	<ul style="list-style-type: none"> 1. <u>Test is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event</u> 2. <u>Test is not appropriate for multiple observations per patient</u> 3. <u>Confidence intervals and/or p values not reported</u> 4. <u>Comparative treatment effects not calculated</u>

CGM: continuous glucose monitoring; CI: confidence interval; RCT: randomized controlled trial.

Pregnant Women

As discussed in the section on CGM in pregnant women, 2 RCTs have evaluated intermittent CGM glucose monitoring in pregnant women with type 1 and type 2 diabetes. Most women had type 1 diabetes in both trials. There were 25 (35%) women with type 2 diabetes in Murphy et al (2008) and 31 (20%) with type 2 diabetes in Secher et al (2013). Results for women with type 2 diabetes were not reported in Murphy. Secher reported that 5 (17%) women with type 2 diabetes experienced 15 severe hypoglycemic events, with no difference between the groups; other analyses were not stratified by diabetes type.

Section Summary: Type 2 Diabetes for Long- and Short-Term Glucose Monitoring

Most RCTs of CGM in patients with type 2 trials found statistically significant benefits of CGM regarding glycemic control. However, the degree of HbA1c reduction and the difference in HbA1c reduction between groups might not be clinically significant. Also, the variability among interventions makes it difficult to identify an optimal approach to CGM use; the studies used a combination of intermittent and continuous monitoring with a review of data in real-time or at study visits only. Only the DIAMOND trial (N=158) used continuous, real-time CGM. Entry criteria regarding the use of insulin and HbA1c levels also varied across studies, and a subgroup of type 2 diabetes patients who might benefit has not been identified. Moreover, studies of CGM in patients with type 2 diabetes generally do not address the clinically important issues of severe hypoglycemia and diabetic complications. The DIAMOND trial reported a larger reduction in change in HbA1c level from baseline to 24 weeks with CGM and a larger proportion of patients with a relative reduction in the HbA1c level of 10% or higher at 24 weeks favoring CGM but no differences in quality of life. There were no events of severe hypoglycemia or diabetic ketoacidosis in either group.

Two trials of CGM have enrolled pregnant women with type 2 diabetes, but the total number of women with type 2 diabetes included in both trials is only 58. One study reported a difference in HbA1c levels at 36 weeks, and the proportion of infants that were large for gestational age (>90th percentile) favored CGM while the second study did not. Neither study reported analyses stratified by diabetes type.

Pregnant Women with Gestational Diabetes

One trial of glucose monitoring in women with gestational diabetes has been published. Trial design, results, and gaps are shown in Tables 18 to 21. In an RCT, Wei et al (2016) evaluated the use of CGM in 120 women with gestational diabetes at 24 to 28 weeks. Patients were allocated to prenatal care plus CGM (n=58) or SMBG (n=62). The CGM sensors were reportedly inserted for 48 to 72 hours on weekdays; it is not clear whether the readings were available in real-time. The investigators assessed a number of end points and did not specify primary outcomes; a significance level of p less than 0.05 was used for all outcomes. The groups did not differ significantly in a change in most outcomes, including a change in maternal HbA1c levels, rates of preterm delivery before the 35th gestational week, cesarean delivery rates, proportions of large-for-gestational age infants, or rates of neonatal hypoglycemia. Women in the CGM group gained significantly less weight than those in the SMBG group.

Table 18. Key RCT Characteristics for CGM in Pregnant Women with Gestational Diabetes

<u>Author</u>	<u>Countries</u>	<u>Sites</u>	<u>Dates</u>	<u>Participants</u>	<u>Interventions</u>	
					<u>Active</u>	<u>Comparator</u>
<u>Wei et al (2016)</u>	<u>China</u>	<u>1</u>	<u>2011-2012</u>	<u>Pregnant women with gestational diabetes diagnosed between 24 and 28 wk' gestation; mean HbA_{1c} level, 5.8%; mean age, 30 y</u>	<u>CGM (48- 721 on weekdays) (n=51)</u>	<u>SMBG (n=55)</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c. RCT: randomized controlled trial; SMBG: self-monitored blood glucose.

Table 19. RCT Outcomes for CGM in Pregnant Women with Gestational Diabetes

<u>Study</u>	<u>Large-for-Gestational Age</u>	<u>Infant</u>		<u>Caesarean Section</u>	<u>Maternal</u>	
		<u>Gestational Age at Delivery, wk</u>	<u>Severe Hypoglycemia</u>		<u>HbA_{1c} Levels at 36 Wk' Gestation^a</u>	<u>Severe Hypoglycemia</u>
<u>Wei et al (2016)</u>						
<u>N</u>	<u>106</u>	<u>106</u>	<u>106</u>	<u>106</u>		<u>NR</u>
<u>CGM</u>	<u>18 (35%)</u>	<u>Mean, 37.4</u>	<u>4 (8%)</u>	<u>31 (60%)</u>	<u>Mean, 5.5%</u>	
<u>Control</u>	<u>29 (53%)</u>	<u>Mean, 37.5</u>	<u>7 (13%)</u>	<u>38 (69%)</u>	<u>Mean, 5.6%</u>	
<u>TE (95% CI)</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	<u>NR</u>	
<u>p</u>	<u>0.07</u>	<u>0.92</u>	<u>0.41</u>	<u>0.37</u>	<u>0.09</u>	

Table 20. Relevance Gaps of RCTs for CGM in Pregnant Women with Gestational Diabetes

<u>Study</u>	<u>Population</u>	<u>Intervention</u>	<u>Comparator</u>	<u>Outcomes</u>	<u>Follow-Up</u>
<u>Wei et al (2016)</u>	<u>4. Study population had relatively low HbA_{1c} level</u>	<u>4. Compliance with CGM not reported</u>	<u>None noted</u>	<u>None noted</u>	<u>None noted</u>
<u>Key</u>	<u>1. Intended use population unclear</u> <u>2. Clinical context for treatment is unclear</u> <u>3. Study population unclear</u> <u>4. Study population not representative of intended use</u> <u>5. Study population is subpopulation of intended use</u>	<u>1. Not clearly defined</u> <u>2. Version used unclear</u> <u>3. Delivery not similar intensity as comparator</u> <u>4. Not delivered effectively</u>	<u>1. Not clearly defined</u> <u>2. Not standard or optimal</u> <u>3. Delivery not similar intensity as intervention</u> <u>4. Not delivered effectively</u>	<u>1. Key health outcomes not addressed</u> <u>2. Physiologic measures, not validated surrogates</u> <u>3. Not CONSORT reporting of harms measurements</u> <u>5. Clinically significant difference not prespecified</u> <u>6. Clinically significant difference not supported</u>	<u>1. Not sufficient duration for benefits</u> <u>2. Not sufficient duration for harms</u>

CGM: continuous glucose monitoring; HbA_{1c}: hemoglobin A1c; RCT: randomized controlled trial.

Table 21. Study Design and Conduct Gaps of RCTs for CGM in Pregnant Women with Gestational Diabetes

<u>Study</u>	<u>Allocation</u>	<u>Blinding</u>	<u>Selective Reporting</u>	<u>Follow-Up</u>	<u>Power</u>	<u>Statistical</u>
<u>Wei et al (2016)</u>	<u>3. Not reported</u>	<u>1. Not blinded; chance of bias in clinical management</u>	<u>1. Registration not reported</u>	<u>5. Exclusions not well justified</u>	<u>1. No power calculations reported; primary outcome not specified</u>	<u>3, 4. Treatment effects and CIs not calculated</u>
Key	<ul style="list-style-type: none"> <u>1. Participants not randomly allocated</u> <u>2. Allocation not concealed</u> <u>3. Allocation concealment unclear</u> <u>4. Inadequate control for selection bias</u> 	<ul style="list-style-type: none"> <u>1. Not blinded to treatment assignment</u> <u>2. Not blinded outcome assessment</u> <u>3. Outcome assessed by treating physician</u> 	<ul style="list-style-type: none"> <u>1. Not registered</u> <u>2. Evidence of selective reporting</u> <u>3. Evidence of selective publication</u> 	<ul style="list-style-type: none"> <u>1. High loss to follow-up or missing data</u> <u>2. Inadequate handling of missing data</u> <u>3. High number of crossovers</u> <u>4. Inadequate handling of crossovers</u> <u>5. Inappropriate exclusions</u> <u>6. Not intent to treat analysis (per protocol for noninferiority trials)</u> 	<ul style="list-style-type: none"> <u>1. Power calculations not reported</u> <u>2. Power not calculated for primary outcome</u> <u>3. Power not based on clinically important difference</u> 	<ul style="list-style-type: none"> <u>1. Test is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event</u> <u>2. Test is not appropriate for multiple observations per patient</u> <u>3. Confidence intervals and/or p values not reported</u> <u>4. Comparative treatment effects not calculated</u>

CGM: continuous glucose monitoring; RCT: randomized controlled trial.

Section Summary: Pregnant Women With Gestational Diabetes

The RCT in women with gestational diabetes was conducted in China with the intervention starting in the 2nd or 3rd trimester and mean baseline HbA1c level less than 6.0%. The type of CGM monitoring was unclear. Trial reporting was incomplete; however, there were no differences between groups for most reported outcomes.

Summary of Evidence

Type 1 Diabetes

For individuals who have type 1 diabetes who are willing and able to use the device, and have adequate medical supervision, who receive long-term CGM, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. Systematic reviews have generally found that at least in the short-term, long-term CGM resulted in significantly improved glycemic control for adults and children with type 1 diabetes, particularly highly compliant patients. A 2017 individual patient data analysis, pooling data from 11 RCTs, found that reductions in HbA1c levels were significantly greater with real-time CGM than with a control intervention. Two RCTs in patients who used multiple daily insulin injections and were highly compliant with CGM devices during run-in phases found that CGM was associated with a larger reduction in HbA1c levels than previous studies. One of the 2 RCTs prespecified hypoglycemia-related outcomes and reported that time spent in hypoglycemia was significantly less in the CGM group. One RCT in pregnant women with type 1 diabetes, which compares real-time CGM with self-monitoring of blood glucose, has also reported a difference in change in HbA1c levels, an increased percentage of time in the recommended glucose control target range, a smaller proportion of infants who were large for gestational age, a smaller proportion of infants who had neonatal intensive care admissions lasting more than 24 hours, a smaller proportion of infants who had neonatal hypoglycemia requiring treatment, and reduced total length of hospital stay all favoring CGM. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcomes.

For individuals who have type 1 diabetes who receive short-term (intermittent) glucose monitoring, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. The evidence for intermittent short-term monitoring on glycemic control is mixed, and there was no definite improvement in HbA1c levels. Studies have not shown an advantage for intermittent glucose monitoring in reducing severe hypoglycemia events, but the number of events reported is generally small and effect estimates imprecise. The evidence is sufficient to determine the effects of the technology on health outcomes.

Type 2 Diabetes

For individuals who have type 2 diabetes who receive long-term, real-time CGM, the evidence includes RCTs. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. Only the DIAMOND RCT (N=158) has used continuous, real-time CGM in type 2 diabetes. Selected patients were highly compliant during a run-in phase. The difference in change in HbA1c levels from baseline to 24 weeks was -0.3% favoring CGM. The difference in the proportion of patients with a relative reduction in HbA1c level by 10% or more was 22% favoring CGM. There were no differences in the proportions of patients with an HbA1c level less

than 7% at week 24. There were no events of severe hypoglycemia or diabetic ketoacidosis in either group. The treatment groups did not differ in any of the quality of life measures. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have type 2 diabetes who receive short-term, intermittent CGM, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. Systematic reviews of 3 to 4 RCTs have found statistically significant benefits from CGM regarding glycemic control. However, the degree of HbA1c reduction and the difference in HbA1c reductions between groups may not be clinically significant. Also, the small number of RCTs and variability among interventions make it difficult to identify an optimal approach to CGM or a subgroup of type 2 diabetes patients who might benefit. Moreover, studies of CGM in patients with type 2 diabetes have generally not addressed the clinically important issues of severe hypoglycemia and diabetic complications. Very few pregnant women with type 2 diabetes have been included in RCTs. The evidence is insufficient to determine the effects of the technology on health outcomes.

Gestational Diabetes

For individuals who are pregnant with gestational diabetes who receive long-term (continuous) or short-term (intermittent) glucose monitoring, the evidence includes an RCT. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. In the RCT, the type of CGM was unclear. Trial reporting was incomplete; however, there was no difference between the groups for the majority of the reported outcomes. The evidence is sufficient to determine the effects of the technology on health outcomes

Practice Guidelines and Position Statements

American Association of Clinical Endocrinologists and American College of Endocrinology

In 2016, the American Association of Clinical Endocrinologists and American College of Endocrinology published a consensus statement on outpatient glucose monitoring. Following are their recommendations on CGM:

- Type 1 diabetes, adults: “CGM recommended, especially for patients with history of severe hypoglycemia, hypoglycemia unawareness and to assist in the correction of hyperglycemia in patients not at goal. CGM users must know basics of sensor insertion, calibration and real-time data interpretation.”
- Type 1 diabetes, children: Same as adults, except that more training and follow-up is needed.
- Type 2 diabetes receiving insulin, sulfonylureas or glinides: “Data on CGM in T2DM [type 2 diabetes mellitus] are limited at this time. Trials assessing the use of CGM in T2DM are ongoing.”

UK National Institute for Health and Care Excellence (NICE)

The National Institute for Health and Care Excellence updated its guidance on diagnosis and management of type 1 diabetes in adults in 2016. The guidance stated that real-time CGM should not be offered routinely to adults with type 1 diabetes but that it can be considered in the following:

- “Do not offer real-time continuous glucose monitoring routinely to adults with type 1 diabetes”
- ...“adults with type 1 diabetes who are willing to commit to using it at least 70% of the time and to calibrate it as needed, and who have any of the following despite optimized use of insulin therapy and conventional blood glucose monitoring:
 - More than 1 episode a year of severe hypoglycaemia with no obviously preventable precipitating cause.
 - Complete loss of awareness of hypoglycaemia.
 - Frequent (more than 2 episodes a week) asymptomatic hypoglycaemia that is causing
 - Problems with daily activities.
 - Extreme fear of hypoglycaemia.
 - Hyperglycaemia (HbA1c level of 75 mmol/mol [9%] or higher) that persists despite testing at least 10 times a day. Continue real-time continuous glucose monitoring only if HbA1c can be sustained at or below”

American Diabetes Association

The 2017 American Diabetes Association position statement on diabetes includes the following recommendations on CGM:

- “When used properly, continuous glucose monitoring (CGM) in conjunction with intensive insulin regimens is a useful tool to lower A1C in selected adults (aged ≥ 25 years) with type 1 diabetes. (A)*
- Although the evidence for A1C lowering is less strong in children, teens, and younger adults, CGM may be helpful in these groups. Success correlates with adherence to ongoing use of the device. (B)
- CGM may be a useful tool in those with hypoglycemia unawareness and/or frequent hypoglycemic episodes. (C) The Association also recommended that physicians assess individual readiness prior to prescribing CGM and that education, training and support are needed for optimal CGM device implementation.

*Levels of Evidence: A: Clear evidence from well-conducted, generalizable RCTs that are adequately powered; B: Supportive evidence from well-controlled cohort studies; C: Supportive evidence from poorly controlled or uncontrolled studies; E: Expert consensus or clinical experience.

Endocrine Society

In 2016, the Endocrine Society published a clinical practice guideline developed by a task force that included the following recommendations on continuous glucose monitoring:

- “Real-time continuous glucose monitors in adult outpatients
 - We recommend real-time continuous glucose monitoring (RT-CGM) devices for adult patients with T1DM [type 1 diabetes mellitus] who have A1C levels above target and who are willing and able to use these devices on a nearly daily basis.
 - We recommend RT-CGM devices for adult patients with well-controlled T1DM who are willing and able to use these devices on a nearly daily basis.
- Use of continuous glucose monitoring in adults with type 2 diabetes mellitus

- We suggest short-term, intermittent RT-CGM use in adult patients with T2DM (not on prandial insulin) who have A1C levels $\geq 7\%$ and are willing and able to use the device.”

U.S. Preventive Services Task Force Recommendations

Not applicable.

Key Words:

GlucoWatch[®], wrist glucose monitor, Glucose Biographer, AutoSensor, and GlucoWatch[®] G2[™] Biographer, continuous monitoring of glucose in the interstitial fluid, intermittent monitoring of glucose in the interstitial fluid, Continuous Glucose Monitoring System, CGMS, CGMS[®] System Gold[™], Minimed, MiniMed Paradigm 522 or 722 insulin pumps, MiniMed Paradigm Real-Time Insulin Pump and Continuous Glucose Monitoring System, combined continuous subcutaneous insulin infusion and blood glucose monitoring device, DexCom STS Continuous Glucose Monitoring System, CGMS iPro Recorder, Freestyle Navigator[®] Continuous Glucose Monitoring System, Guardian[®] REAL-Time Continuous Glucose Monitoring System, CGM, Dexcom G5, Abbott[®] Freestyle Libre Flash, Dexcom G6

Approved by Governing Bodies:

Several continuous glucose monitoring systems have been approved by the FDA through the premarket approval process:

Table 1. CGM Systems Approved by the Food and Drug Administration

<u>Device</u>	<u>Manufacturer</u>	<u>Approval</u>	<u>Indications</u>
<u>Continuous Glucose Monitoring System (CGMS[®])</u>	<u>MiniMed</u>	<u>1999</u>	<u>3-d use in physician's office</u>
<u>GlucoWatch G2[®] Biographer</u>		<u>2001</u>	<u>Not available since 2008</u>
<u>Guardian[®]-RT (Real-Time) CGMS</u>	<u>MiniMed</u> <u>(now</u> <u>Medtronic)</u>	<u>2005</u>	
<u>Dexcom[®] STS CGMS system</u>	<u>Dexcom</u>	<u>2006</u>	
<u>Paradigm[®] REAL-Time System (second generation called Paradigm Revel System)</u>	<u>MiniMed</u> <u>(now</u> <u>Medtronic)</u>	<u>2006</u>	<u>Integrates a CGM with a Paradigm insulin pump</u>
<u>FreeStyle Navigator[®] CGM System</u>	<u>Abbott</u>	<u>2008</u>	
<u>Dexcom[®] G4 Platinum</u>	<u>Dexcom</u>	<u>2012</u> <u>2014</u>	<u>Adults >18 y; can be worn for up to 7 d</u>
<u>Dexcom[®] G5 Mobile CGM</u>	<u>Dexcom</u>	<u>2016^a</u>	<u>Expanded to include patients with diabetes 2-17 y</u> <u>Replacement for fingerstick blood glucose testing in patients >2 y. System requires at least 2 daily fingerstick tests for calibration purposes, but additional fingersticks are not necessary because treatment decisions can be made based on device readings</u>
<u>Freestyle Libre[®] Pro Flash Glucose Monitoring System</u>	<u>Abbott</u>	<u>2017</u>	<u>Adults' >18 y. Readings are only made available to patients through consultation with a health care professional. Does not require user calibration with blood glucose values</u>

CGM: continuous glucose monitoring.

a As a supplement to the G4 premarketing approval.

Benefit Application:

Coverage is subject to member's specific benefits. Group specific policy will supersede this policy when applicable.

ITS: Home Policy provisions apply

FEP contracts: FEP does not consider investigational if FDA approved and will be reviewed for medical necessity. Special benefit consideration may apply. Refer to member's benefit plan.

Current Coding:

CPT codes:

- 95249** Ambulatory continuous glucose monitoring of interstitial tissue fluid via a subcutaneous sensor for a minimum of 72 hours; patient-provided equipment, sensor placement, hook-up, calibration of monitor, patient training, and printout of recording (**Effective 01/01/2018**)
- 95250** Ambulatory continuous glucose monitoring of interstitial tissue fluid via a subcutaneous sensor for a minimum of **72 hours**; physician or other qualified health care professional (office) provided equipment, sensor placement, hook-up, calibration of monitor, patient training, removal of sensor, and printout of recording
- 95251** Ambulatory continuous glucose monitoring of interstitial tissue fluid via a subcutaneous sensor for a minimum of **72 hours**; analysis, interpretation and report
- 99091** Collection and interpretation of physiologic data (e.g., ECG, blood pressure, glucose monitoring) digitally stored and/or transmitted by the patient and/or caregiver to the physician or other qualified health care professional, qualified by education, training, licensure/regulation (when applicable) requiring a minimum of 30 minutes of time

HCPCS:

- A9276** Sensor; invasive (e.g., subcutaneous), disposable, for use with interstitial continuous glucose monitoring system, one unit = 1 day supply
- A9277** Transmitter; external, for use with interstitial continuous glucose monitoring system
- A9278** Receiver (monitor); external, for use with interstitial continuous glucose monitoring system
- A9999** Miscellaneous DME supply, accessory, and/or service component of another HCPCS code
- E1399** Durable medical equipment, miscellaneous
- K0553** Supply allowance for therapeutic continuous glucose monitor (CGM) system, includes all supplies and accessories, 1 month supply = 1 unit of service. (**Effective 07/01/2017**)

- K0554** Receiver (monitor), dedicated, for use with therapeutic continuous glucose monitor system. (Effective 07/01/2017)
- S1030** Continuous noninvasive glucose monitoring device purchase (for physician interpretation of data, use CPT code)
- S1031** Continuous noninvasive glucose monitoring device, rental, including sensor, sensor replacement, and download to monitor (for physician interpretation of data, use CPT code)

Continuous glucose monitoring (CGM) systems measure glucose in interstitial fluid, rather than capillary blood. Because they do not measure blood glucose, different HCPCS and CPT coding are used for these systems and supplies (HCPCS codes **A9276-A9278** and **K0553-K0554**).

References:

1. Agrawal P, Zhong A, Welsh JB, et al. Retrospective analysis of the real-world use of the threshold suspend feature of sensor-augmented insulin pumps. *Diabetes Technol Ther.* May 2015; 17(5):316-319.
2. American Diabetes Association. 7. Approaches to glycemic treatment. *Diabetes Care.* Jan 2015; 38 (suppl: S41-48).
3. American Diabetes Association. Standards of medical care in diabetes—2010. *Diabetes Care* 2010; 33(suppl 1):S11-61.
4. American Diabetes A. Standards of medical care in diabetes--2013. *Diabetes Care* 2013; 36 Suppl 1:S11-66.
5. American Diabetes Association. Standards in Medical Care in Diabetes, 2014. 2014; www.care.diabetesjournals.org/content/37/Supplement_1/S14.full.pdf+html Accessed June 17, 2016.
6. American Diabetes Association (ADA). 6. Glycemic Targets. *Diabetes Care.* Jan 2017; 40(Suppl 1):S48-S56.
7. American DA. Executive summary: standards of medical care in diabetes—2011. *Diabetes Care* 2011; 34(Suppl 1):S4-S10.
8. Bailey KJ, Little JP, Jung ME. Self-monitoring using continuous glucose monitors with real-time feedback improves exercise adherence in individuals with impaired blood glucose: a pilot study. *Diabetes Technol Ther.* Mar 2016; 18(3):185-193.
9. Battelino T, Conget I, Olsen B et al. The use and efficacy of continuous glucose monitoring in type 1 diabetes treated with insulin pump therapy: a randomized controlled trial *Diabetologia* 2012; 55(12):3155-3162.
10. Beck RW, Riddlesworth TD, Ruedy K, et al. Continuous glucose monitoring versus usual care in patients with type 2 diabetes receiving multiple daily insulin injections: a randomized trial. *Ann Intern Med.* Sep 19 2017; 167(6):365-374.
11. Beck RW, Riddlesworth T, Ruedy K, et al. Effect of continuous glucose monitoring on glycemic control in adults with type 1 diabetes using insulin injections: The DIAMOND randomized clinical trial. *Jama.* Jan 24 2017; 317(4):371-378.
12. Benkhadra K, Alahdab F, Tamhane S, et al. Real-time continuous glucose monitoring in type 1 diabetes: a systematic review and individual patient data meta-analysis. *Clin Endocrinol (Oxf).* Mar 2017; 86(3):354-360.

13. Bergenstal RM, Klonoff DC, Garg SK et al. Threshold-based insulin-pump interruption for reduction of hypoglycemia. *N Engl J Med* 2013; 369(3):224-32.
14. Bloomgarden ZT. American Diabetes Association 60th Scientific Sessions 2000. Glucose tolerance, diabetes and cancer, glycemic control, monitoring and related topics. *Diabetes Care*, April 2001, 24(4): 779-784.
15. Blue Cross Blue Shield Association Technology Evaluation Criteria (TEC) Assessment. Use of intermittent or continuous interstitial fluid glucose monitoring in patients with diabetes mellitus. *TEC Assessments 2003; Volume 18, Tab 16.*
16. Bode BW, et al. What's ahead in glucose monitoring? New techniques hold promise for improved ease and accuracy. *Postgrad Med*, April 2001, 109(4): 41-4, 47-9.
17. Buckingham, B. Continuous glucose monitoring in children with type 1 diabetes. *J Pediatr* 2007; 151:388-93.
18. Burge MR, Mitchell S, Sawyer A, Schade DS. Continuous glucose monitoring: the future of diabetes management. *Diabetes Spectrum* 2008; 21(2):112-19.
19. Chetty VT, Almulla A, Oduyungbo A et al. The effect of continuous subcutaneous glucose monitoring (CGMS) versus intermittent whole blood finger-stick glucose monitoring (SBGM) on hemoglobin A1C (HBA1c) levels in Type I diabetic patients: a systematic review. *Diabetes Res Clin Pract* 2008; 81(1):79-87.
20. Chico A, Vidal-Rios, P, Subira M, Novials A. The continuous glucose monitoring system is useful for detecting unrecognized hypoglycemia in patients with type 1 and type 2 diabetes but is not better than frequent capillary glucose measurements for improving metabolic control. *Diabetes Care* 2003; 26(4):1153-57.
21. Deiss D, Hartmann R, Schmidt J, Kordonouri O. Results of a randomised controlled cross-over trial on the effect of continuous subcutaneous glucose monitoring (CGMS) on glycaemic control in children and adolescents with type 1 diabetes. *Exp Clin Endocrinol Diabetes*. 2006 114: 63-67.
22. Deiss D, Bolinder J, Riveline JP, et al. Improved glycemic control in poorly controlled patients with type 1 diabetes using real-time continuous glucose monitoring. *Diabetes Care* 2006; 29(12):2730-32.
23. Effect of intensive therapy on the microvascular complications of type 1 diabetes mellitus. *Jama*. May 15 2002; 287(19):2563-2569.
24. Erhardt NM, Chellapa M, Walker MS et al. The effect of real-time continuous glucose monitoring on glycemic control in patients with type 2 diabetes mellitus. *J Diabetes Sci Technol* 2011; 5(3):668-75.
25. Feig DS, Donovan LE, Corcoy R, et al. Continuous glucose monitoring in pregnant women with type 1 diabetes (CONCEPTT): a multicentre international randomised controlled trial. *Lancet*. Nov 25 2017; 390(10110):2347-2359.
26. Floyd B, Chandra P, Hall S et al. Comparative analysis of the efficacy of continuous glucose monitoring and self-monitoring of blood glucose in type 1 diabetes mellitus. *J Diabetes Sci Technol* 2012; 6(5):1094-1102.
27. Food and Drug Administration (FDA). Summary of Safety and Effectiveness (SSED): Dexcom G5 Mobile Continuous Glucose Monitoring System. 2016; www.accessdata.fda.gov/cdrh_docs/pdf12/P120005S041b.pdf. Accessed May 31, 2017.
28. Food and Drug Administration (FDA). News Release: Freestyle Libre Flash Glucose Monitoring System. 2017;

<https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm577890.htm>.
Accessed January 11, 2018.

29. Gandhi GY, Kovalaske M, Kudva Y et al. Efficacy of continuous glucose monitoring in improved glycemic control and reducing hypoglycemia: a systematic review and meta-analysis of randomized trials. *J Diabetes Sci Technol* 2011; 5(4):952-65.
30. Garg SK, et al. Correlation of fingerstick blood glucose measurements with GlucoWatch biographer glucose results in young subjects with type 1 diabetes. *Diabetes Care*, October 1999, 22(10): 1708-14.
31. Garg SK, et al. Improved glucose excursions using an implantable real-time continuous glucose sensor in adults with type 1 diabetes. *Diabetes Care*, March 2004, Vol. 27, No. 3.
32. Garg SK, et al. Improvement in glycemic excursions with a transcutaneous Real Time continuous glucose sensor. *Diabetes Care*, January 2006.
33. Garg S, Brazg RL, Bailey TS et al. Reduction in duration of hypoglycemia by automatic suspension of insulin delivery: the in-clinic ASPORE study. *Diab Technol Ther* 2012; 14(3):205-9.
34. Gehlert RR, Dogbey GY, Schwartz FL, et al. Hypoglycemia in type 2 diabetes--more common than you think: a continuous glucose monitoring study. *J Diabetes Sci Technol*. Sep 2015; 9(5):999-1005.
35. Gold AE, MacLeod KM, Frier BM. Frequency of severe hypoglycemia in patients with type I diabetes with impaired awareness of hypoglycemia. *Diabetes Care*. Jul 1994; 17(7):697-703.
36. Guerci B, Floriot M, Böhme P, et al. Clinical performance of CGMS in type 1 diabetic patients treated by continuous subcutaneous insulin infusion using insulin analogs. *Diabetes Care* 2003; 26(3):582-89.
37. Jovanovic L, et al. A randomized controlled study of a transcutaneous, Real Time continuous glucose sensor demonstrates improvement in glycemic control. The 65th Scientific Sessions of the American Diabetes Association, June 2005.
38. Jovanovic L, et al. Results from an unblinded study of a short-term continuous glucose sensor in subjects with type 1 diabetes. The 65th Scientific Sessions of the American Diabetes Association, June 2005.
39. Juvenile Diabetes Research Foundation Continuous Glucose Monitoring Study Group. Continuous glucose monitoring and intensive treatment of type 1 diabetes. *N Engl J Med* 2008; 359(14):1464-1476.
40. Juvenile Diabetes Research Foundation Continuous glucose Monitoring Study Group. The effect of continuous glucose monitoring in well-controlled type 1 diabetes. *Diabetes Care* 2009; 32(8):1378-1383.
41. Juvenile Diabetes Research Foundation Continuous Glucose Monitoring Study Group. Effectiveness of continuous glucose monitoring in a clinical care environment. *Diabetes Care* 2010; 33(1):17-22.
42. Kaufman FR, et al. A pilot study of the continuous glucose monitoring system. *Diabetes care* 2001; 24(12):2030-34.
43. Klonoff DC. Continuous glucose monitoring: roadmap for 21st century diabetes therapy. *Diabetes Care* 2005; 28(5):1231-39.
44. Klonoff DC, Buckingham B, Christiansen JS et al. Continuous glucose monitoring: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011; 96(10):2968-79.

45. Koschinsky T and Heinemann L. Sensors for glucose monitoring: Technical and clinical aspects. *Diabetes Metab Res Rev*, March-April 2001, 17(2): 113-23.
46. Langendam M, Luijf YM, Hooft L et al. Continuous glucose monitoring systems for type 1 diabetes mellitus. *Cochran Database Syst Rev* 2012; 1:CD0081010.
47. Lind M, Polonsky W, Hirsch IB, et al. Continuous glucose monitoring vs conventional therapy for glycemic control in adults with type 1 diabetes treated with multiple daily insulin injections: The GOLD randomized clinical trial. *Jama*. Jan 24 2017; 317(4):379-387.
48. Little SA, Leelarathna L, Walkinshaw E, et al. Recovery of hypoglycemia awareness in long-standing type 1 diabetes: a multicenter 2 x 2 factorial randomized controlled trial comparing insulin pump with multiple daily injections and continuous with conventional glucose self-monitoring (HypoCOMPASS). *Diabetes Care*. Aug 2014; 37(8):2114-2122.
49. Ly TT, Nicholas JA, Retterath A, et al. Effect of sensor-augmented insulin pump therapy and automated insulin suspension vs standard insulin pump therapy on hypoglycemia in patients with type 1 diabetes: a randomized clinical trial. *JAMA*. Sep 25 2013; 310(12):1240-1247.
50. Management of Diabetes in Pregnancy: Standards of Medical Care in Diabetes-2018. *Diabetes Care*. Jan 2018; 41(Suppl 1):S137-s143.
51. Mauras N, Beck R, Xing D et al. A randomized clinical trial to assess the efficacy and safety of real-time continuous glucose monitoring in the management of type 1 diabetes in young children aged 4 to <10 years. *Diabetes Care* 2012; 35(2):204-210.
52. McGowan K, Thomas W, Moran A. Spurious reporting of nocturnal hypoglycemia by CGMS in patients with tightly controlled type 1 diabetes. *Diabetes care* 2002; 25(9):1499-1503.
53. Murphy HR, Rayman G, Lewis K et al. Effectiveness of continuous glucose monitoring in pregnant women with diabetes: randomised clinical trial. *BMJ* 2008; 337:a1680.
54. National Center for Health and Care Excellence (NICE). Type 1 diabetes in adults: diagnosis and management. www.nice.org.uk/guidance/ng17?unlid=382286372016220232952. Accessed June 10, 2016.
55. National Institute for Health and Care Excellence. Integrated sensor-augmented pump therapy systems for managing blood glucose levels in type 1 diabetes (the MiniMed Paradigm Veo system and the Vibe and G4 PLATINUM CGM system). Diagnostics guidance [DG21]. Feb 2016. <https://www.nice.org.uk/guidance/dg21/chapter/1-Recommendations>. Accessed October 21, 2016.
56. Newman SP, Hurel SJ, Cooke D, et al. A randomized control trial of continuous glucose monitoring devices on HbA1c – The MITRE study. *American Diabetes Association* 2007.
57. Newman SP, Cooke D, Casbard A et al. A randomized controlled trial to compare minimally invasive glucose monitoring devices with conventional monitoring in the management of insulin-treated diabetes mellitus (MITRE). *Health Technol asses* 2009; 13(28): iii-iv, 1-194.
58. Pazos-Couselo M, Garcia-Lopez JM, Gonzalez-Rodriguez M, et al. High incidence of hypoglycemia in stable insulin-treated type 2 diabetes mellitus: continuous glucose monitoring vs. self-monitored blood glucose. *Observational prospective study. Can J Diabetes*. Oct 2015; 39(5):428-433.
59. Peters AL, Ahmann AJ, Battelino T, et al. Diabetes technology-continuous subcutaneous insulin infusion therapy and continuous glucose monitoring in adults: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. Nov 2016; 101(11):3922-3937.

60. Pitzer KR, et al. Detection of hypoglycemia with the GlucoWatch biographer. *Diabetes Care*, May 2001, 24(5): 881-885.
61. Polonsky WH, Hessler D, Ruedy KJ, et al. The impact of continuous glucose monitoring on markers of quality of life in adults with type 1 diabetes: further findings from the DIAMOND randomized clinical trial. *Diabetes Care*. Jun 2017; 40(6):736-741.
62. Poolsup N, Suksomboon N, Kyaw AM. Systematic review and meta-analysis of the effectiveness of continuous glucose monitoring (CGM) on glucose control in diabetes. *Diabetol Metab Syndr* 2013; 5(1):39.
63. Raccach D, Sulmont V, Reznik Y et al. Incremental value of continuous glucose monitoring when starting pump therapy in patients with poorly controlled type 1 diabetes: the RealTrend study. *Diabetes Care* 2009; 32(12):2245-50.
64. Phillip M, Battelino T, Atlas E et al. Nocturnal glucose control with an artificial pancreas at a diabetes camp. *N Engl J Med* 2013; 368(9):824-33.
65. Robert JJ. Continuous monitoring of blood glucose. *Hormone Research*, January 2002; 57 Suppl 1: 81-4. (Abstract)
66. Sato J, Kanazawa A, Ikeda F, et al. Effect of treatment guidance using a retrospective continuous glucose monitoring system on glycaemic control in outpatients with type 2 diabetes mellitus: A randomized controlled trial. *J Int Med Res*. Feb 2016; 44(1):109-121.
67. Secher AL, Ringholm L, Andersen HU et al. The Effect of Real-Time Continuous Glucose Monitoring in Pregnant Women with Diabetes: a randomized controlled trial. *Diabetes care* 2013 Jul 2013; 36(7):1877-1883.
68. Sequeira PA, Montoya L, Ruelas V, et al. Continuous glucose monitoring pilot in low-income type 1 diabetes patients. *Diabetes Technol Ther*. Oct 2013; 15(10):855-858.
69. Silverstein JH and Rosenbloom AL. New developments in type 1 (insulin-dependent) diabetes. *Clin Pediatr (Phila)*, May 2000, 39(5): 257-66.
70. Tamada JA, et al. Non-Invasive glucose monitoring: Comprehensive clinical results. *JAMA*, November 1999, 282(19): 1839-1844.
71. The Diabetes Research in Children Network (DirecNet) Study Group. Accuracy of the GlucoWatch G2 Biograpphar and the continuous glucose monitoring system during hypoglycemia. *Diabetes Care* 2004; 27(3):722-26.
72. Tierney MJ, et al. The GlucoWatch biographer: A frequent automatic and noninvasive glucose monitor. *Ann Med*, December 2000, 32(9): 632-41.
73. van Beers CA, DeVries JH, Kleijer SJ, et al. Continuous glucose monitoring for patients with type 1 diabetes and impaired awareness of hypoglycemia (IN CONTROL): a randomized, open-label, crossover trial. *Lancet Diabetes Endocrinol*. Nov 2016; 4(11):893-902.
74. Vigersky RA, Fonda SJ, Chellappa M et al. Short-and long-term effects of real-time continuous glucose monitoring in patients with type 2 diabetes. *Diabetes Care* 2012; 35(1):32-38.
75. Voormolen DN, Devries JH, Frax A et al. Effectiveness of continuous glucose monitoring during diabetic pregnancy (GlucoMOMS trial); a randomised controlled trial. *BMCPrnancy Childbirth* 2012; 12(1):164.
76. Voormolen DN, Devries JH, Evers IM et al. The efficacy and effectiveness of continuous glucose monitoring during pregnancy: a systematic review. *Obstet Gynecol Surv* 2013; 68(11):753-63.

77. Wainstein J, et al. Insulin pump therapy vs. multiple daily injections in obese Type 2 diabetic patients. *Diabetes Medicine* 2005; 22(8): 1037-1046.
78. Wei Q, Sun Z, Yang Y, et al. Effect of a CGMS and SMBG on maternal and neonatal outcomes in gestational diabetes mellitus: a randomized controlled trial. *Sci Rep.* 2016; 6:19920.
79. Wentholt IME, Hoekstra JBL, Devries JH. Continuous glucose monitors: the long-awaited watch dogs? *Diabetes Technology & Therapeutics* 2007; 9(5):399-09.
80. Wilson DM, Beck RW, Tamborlane WV, et al. The accuracy of the FreeStyle Navigotor continuous glucose monitoring system in children with type 1 diabetes. *Diabetes Care* 2007; 30(1):59-64.
81. Wojcichowski P, Rys P, Lipowska A et al. Efficacy and safety comparison of continuous glucose monitoring and self-monitoring of blood glucose in type 1 diabetes. *Pool Arch Med Wewn* 2011; 121(10):333-343.
82. Wolpert HA. The nuts and bolts of achieving end points with real-time continuous glucose monitoring. *Diabetes Care* 2008; 31(suppl 2):S146-9.
83. Yeoh E, Choudhary P, Nwokolo M, et al. Interventions that restore awareness of hypoglycemia in adults with type 1 diabetes: a systematic review and meta-analysis. *Diabetes Care.* Aug 2015; 38(8):1592-1609.
84. Zavalkoff Samara R and Polychronakos Constantin. Evaluation of conventional blood glucose monitoring as an indicator of integrated glucose values using a continuous subcutaneous sensor. *Diabetes Care*, September 2002, Vol. 25(9), pp. 1603-1606.

Policy History:

Medical Policy Administration Committee, March 2002
 Available for Comment April 15-May 29, 2002
 Medical Policy Group, March 2003 (1)
 Medical Policy Administration Committee, March 2003
 Available for comment April 1-May 16, 2003
 Medical Policy Group, March 2004
 Medical Policy Administration Committee, April 2004
 Available for comment April 6-May 20, 2004
 Medical Policy Group, March 2005 (1)
 Medical Policy Group, March 2006 (1)
 Medical Policy Group, June 2006 (2)
 Medical Policy Administration Committee, June 2006
 Available for comment July 7-August 21, 2006
 Medical Policy Group, January 2007 (3)
 Medical Policy Administration Committee, February 2007
 Available for comment February 10-March 26, 2007
 Medical Policy Group, July 2007 (1)
 Medical Policy Group, October 2007 (2)
 Medical Policy Administration Committee, October 2007
 Available for comment October 20-December 3, 2007
 Medical Policy Group, June 2008 (2)
 Medica Review Committee, June 2008

Medical Policy Administration Committee, July 2008
 Available for comment August 7-September 22, 2008
 Medical Policy Group, November 2008 (2)
 Medical Policy Administration Committee, December 2008
 Medical Policy Group, January 2009 (2)
 Medical Policy Group, February 2009 (2)
 Medical Policy Panel, March 2010
 Medical Policy Group, May 2010(2)
 Medical Policy Group, December 2012 (3): 2013 Coding Updates: Verbiage changes to Code 99091.
 Medical Policy Panel, March 2013
 Medical Policy Group, May 2013 (2): Policy statement added that artificial pancreases are considered investigational. Definition of poorly controlled Type 1 diabetes added. Definition of best practices added. Description, Key Points, Key Words, and References updated to support Policy Statements. Policy Statements prior to June 28, 2008 removed.
 Medical Policy Administration Committee, June 2013
 Available for Comment May 30 through July 13, 2013
 Medical Policy Group, April 2014 (5): Policy statement added for coverage of sensor-augmented insulin pump therapy with the low glucose threshold suspend feature; Sensor-augmented insulin pump therapy with the low glucose threshold suspend feature is considered investigational in children younger than 16 years; Replacement or upgrade of properly functioning equipment, even if warranty has expired, does not meet Blue Cross and Blue Shield of Alabama’s medical criteria for coverage. Removed closed-loop monitoring device with LGS feature from being considered investigational. Description, Key Points, Key Words, Practice Guidelines & Position Statements, Approved Governing Bodies, and References updated to support Policy Statements.
 Medical Policy Administration Committee, April 2014
 Available for comment April 4 through May 19, 2014
 Medical Policy Group, May 2014 (5): Added new codes S1034, S1035, S1036, and S1037 effective 7/1/2014.
 Medical Policy Group, January 2015 (6): Updates – Key Points and References; no change to policy statement.
 Medical Policy Panel, June 2016
 Medical Policy Group, July 2016 (6): Updated: Description, Key Points, Summary, U.S. Preventive Services Task Force Recommendations, Key Words, Practice Guidelines and Position Statements and References; No change to policy statement.
 Medical Policy Panel, November 2016
 Medical Policy Group, December 2016 (6): Updated Description, Key Points, Key Words, Governing Bodies, Summary, Practice Guidelines and References.
 Medical Policy Group, February 2017 (6): Removed Artificial Pancreas information from Description, Policy statement, Key Points, Key Words, Practice Guidelines, Governing Bodies, References and Coding. Moved to policy #636, Artificial Pancreas Device Systems.
 Medical Policy Group, June 2017: Quarterly Coding update. Added HCPCs K0553 and K0554 to Current Coding.
 Medical Policy Panel, July 2017
 Medical Policy Group, July 2017 (6): Updates to Description, Key Points, Practice Guidelines, Governing Bodies, Key Words, Coding and References.

Medical Policy Group, September 2017 (6): Updated policy statement bullet to include “or impaired awareness”.

Medical Policy Group, November 2017 (6): Updated coding section to address system limitations of providers for billing the Dexcom G5 CGM.

Medical Policy Group, December 2017: Annual Coding Update 2018. Added new CPT code 95249 effective 1/1/18 to Current Coding; updated description for revised codes 95250 and 95251.

Medical Policy Group, December 2017 (6): Removed coding instructions from Coding Section.

Medical Policy Group, January 2018 (6): Updated Governing Bodies, Key Words and References to include the Abbott Freestyle Libre CGM. No change to policy statement.

Medical Policy Group, January 2018 (6): Added verbiage to policy section: Coverage for non-medical items, even when the items may be used to serve a medical purpose, such as smart devices (smart phones, tablets, personal computers, etc.) are non-covered. This includes smart devices used in conjunction with Continuous Glucose Monitors.

Medical Policy Panel, March 2018

Medical Policy Group, May 2018 (6): Updates to Description, Key Points, Governing Bodies, Key Words and References.

This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member’s plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.

This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield’s administration of plan contracts.