2025 COMMON HCC ICD-10 CODES



When documenting chronic conditions, remember: 1) Coding is based on clear documentation that includes a diagnostic statement and ongoing treatment plan. Any condition that is taken into account or affects patient care, treatment or management at the time of the encounter should be documented and coded. 2) The encounter note should be a face-to-face visit and be complete, legible, concise, and contain the provider signature with credentials. 3) Avoid the use of uncertain diagnoses such as "suggestive of," "suspected," "consistent with" or "probable," and code to the highest level of certainty for the encounter/visit.

| Diabetes Mellitus (DM) | TYPE 1 | TYPE 2 | Alzheimer's disease unspecified |
|---|---------------------------------|------------------|---|
| Diabetes mellitus without complications | E10.9 | E11.9 | Alzheimer's disease: Must be specifically conf |
| Combination Codes: Utilize combination codes that link the complications disease. If there are multiple complications of diabetes mellitus then be sure each of the diabetes mellitus combination codes. | | | that the Alzheimer's code be paired with the add Alzheimer's. The physician does not have to me Epilepsy, unspecified, not intractable, witho |
| | C10 CE | F11 CE | Unspecified convulsions |
| Diabetes mellitus with hyperglycemia | E10.65 | E11.65 | <u> </u> |
| Hyperglycemia: When hyperglycemia, poorly controlled but of control is documented, diabetes mellitus with hyp. The word uncontrolled can mean either hypo- or hypergonsufficient documentation to code E10.65/E11.65. | erglycemia sho | ould be coded. | Do not assign R56.9 when a patient has had a seizure disorder or recurrent seizures are preserved. Cardiovascular Conditions |
| Diabetes mellitus with diabetic neuropathy, unspecified | E10.40 | E11.40 | Hypertensive heart disease with heart failur |
| Diabetes mellitus with polyneuropathy | E10.42 | E11.42 | 31 |
| Jabetes menitus with polyneuropathy | E10.42 | E11.42 | Use additional code to identify type of hear |
| Diabetes Mellitus with Kidney Complications | TYPE 1 | TYPE 2 | Hypertensive heart disease without heart failure diagnosis and a heart disease code from the rar |
| Diabetes mellitus with diabetic chronic kidney disease | E10.22 | E11.22 | code is needed other than I11.9. |
| Diabetes mellitus with diabetic nephropathy | E10.21 | E11.21 | Hypertensive chronic kidney disease with s |
| Diabetic chronic kidney disease (CKD) and nephro | pathy: When | | disease or end stage renal disease |
| nephropathy and CKD are documented, code diabetic | CKD, not neph | ropathy. | Use additional code to identify stage of CK |
| Jse additional code to also to identify stage of CKD. | N18.1- N18.6 | N18.1- N18.6 | Hypertensive chronic kidney disease with s chronic kidney disease, or unspecified chro |
| Diabetes Mellitus with Ophthalmic Complications | TYPE 1 | TYPE 2 | Use additional code to identify stage of CKI |
| Diabetes mellitus with unspecified diabetic retinopathy | E10.31 | E11.31 | , , |
| Ophthalmic complications: f adequately documented, code specifically as to type and designate right, left, bilateral or unspecified eye. | E10.3- | E11.3- | Hypertensive heart and chronic kidney disea and stage 1 through stage 4 chronic kidney chronic kidney disease |
| ind designate right, left, bilateral of unspecified eye. | | | Use additional code to identify heart failure |
| Diabetes Mellitus with Circulatory Complications | TYPE 1 | TYPE 2 | stage of CKD. |
| Diabetic peripheral angiopathy without gangrene | E10.51 | E11.51 | Use additional code to identify stage of CK |
| Diabetic peripheral angiopathy with gangrene | E10.52 | E11.52 | |
| Diabetes mellitus with foot ulcer | E10.621 | E11.621 | Hypertensive heart and chronic kidney disea |
| Jse additional code to identify site of ulcer. | L97.4-, L97.5 | - L97.4-, L97.5- | and with stage 5 chronic kidney disease or e |
| | | | Use additional code to identify stage of CKI |
| Endocrine, Nutritional and Metabolic Disorders | | CODE | Hypertensive heart and chronic kidney disease |
| Morbid (severe) obesity due to excess calories | | E66.01 | with stage 5 chronic kidney disease or end s |
| Evaluate for morbid obesity in patients with BMI > or a norbid obesity for patients with comorbidities affected | = to 40. Also e d by weight. | valuate for | Use additional code to identify the type of stage of CKD. |
| Mental, Behavioral and Neurodevelopmental | | CODE | Pulmonary hypertension, unspecified |
| <u> </u> | | | Pulmonary hypertension due to left heart d |
| Dementia, unspecified without behavioral disturbance | | F03.90- | Pulmonary hypertension due to lung diseas |
| Dementia, unspecified with behavioral disturbance | | F03.91- | Cor pulmonale (Chronic) |
| Major Depressive Disorder, single episode, moderate | | F32.1 | Paroxysmal atrial fibrillation |
| Major Depressive Disorder, single episode, severe, wit osychotic features | hout | F32.2 | Longstanding persistent atrial fibrillation |
| Major Depressive Disorder, single episode, severe, wit | h nevehotic | F32.3 | Defined as persistent and continuous lastin |
| viajor Depressive Disorder, sirigie episode, severe, wit features | ii psycholic | 1 32.3 | Other persistent atrial fibrillation |
| Major Depressive Disorder, single episode, in full remi | ssion | F32.5 | Code I48.19 when documented as chronic |
| Major Depressive Disorder, recurrent, moderate | | F33.1 | Chronic atrial fibrillation, unspecified |
| Major Depressive Disorder, recurrent, severe, without psyc | | F33.2 | Only code chronic AF when documented by |
| Major Depressive Disorder, recurrent, severe, with psych | | F33.3 | Permanent atrial fibrillation |
| Document the following components of depression: | odo rodialos | . 50.0 | Not to be confused with persistent; assign onl |
| Degree (mild, moderate, severe); Episode (single (partial or full remission); Presence or absence of p | | | Unspecified atrial fibrillation |
| | _ | | Unspecified atrial flutter |
| Anorexia nervosa, unspecified | | F50.00 | Sick sinus syndrome |
| A | | EEO 04 | |

Anorexia nervosa, restricting type

Bulimia nervosa

Anorexia nervosa, binge eating/purging type

| provider signature with credentials. 3) Avoid the use of uncertain diagnoses such est level of certainty for the encounter/visit. | | |
|--|-----------------------------|--|
| Alzheimer's disease unspecified | G30.9 | |
| Alzheimer's disease: Must be specifically confirmed by the physician to c that the Alzheimer's code be paired with the additional F02.8- codes as a r Alzheimer's. The physician does not have to mention dementia to code it. | | |
| Epilepsy, unspecified, not intractable, without status epilepticus | G40.909 | |
| Unspecified convulsions | R56.9 | |
| Do not assign R56.9 when a patient has had a seizure disorder or recurrent seizure disorder or recurrent seizures are present utilize appropriate code f | | |
| Cardiovascular Conditions | CODE | |
| Hypertensive heart disease with heart failure | l11.0 | |
| Use additional code to identify type of heart failure. | 150 | |
| Hypertensive heart disease without heart failure can be coded when there i diagnosis and a heart disease code from the range I51.4-I51.7; I51.89; I5 code is needed other than I11.9. | | |
| Hypertensive chronic kidney disease with stage 5 chronic kidney disease or end stage renal disease | 112.0 | |
| Use additional code to identify stage of CKD. | N18.5-N18.6 | |
| Hypertensive chronic kidney disease with stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease | 112.9 | |
| Use additional code to identify stage of CKD. | N18.1-N18.4, N18.9 | |
| Hypertensive heart and chronic kidney disease with heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease | 113.0 | |
| Use additional code to identify heart failure type and stage of CKD. | I50, N18.1- N18.4, N18.9 | |
| Use additional code to identify stage of CKD. | N18.1-N18.4, N18.9 | |
| Hypertensive heart and chronic kidney disease without heart failure and with stage 5 chronic kidney disease or end stage renal disease | 113.11 | |
| Use additional code to identify stage of CKD. | N18.5, N18.6 | |
| Hypertensive heart and chronic kidney disease with heart failure and with stage 5 chronic kidney disease or end stage kidney disease | 113.2 | |
| Use additional code to identify the type of heart failure and stage of CKD. | I50, N18.5, N18.6 | |
| Pulmonary hypertension, unspecified | 127.20 | |
| Pulmonary hypertension due to left heart disease | 127.22 | |
| Pulmonary hypertension due to lung disease and hypoxia | 127.23 | |
| Cor pulmonale (Chronic) | 127.81 | |
| Paroxysmal atrial fibrillation | 148.0 | |
| Longstanding persistent atrial fibrillation | 148.11 | |
| Defined as persistent and continuous lasting longer than a year. | I . | |
| Other persistent atrial fibrillation | 148.19 | |
| Code I48.19 when documented as chronic persistent AF. | I | |
| Chronic atrial fibrillation, unspecified | 148.20 | |
| Only code chronic AF when documented by the provider. | 140.04 | |
| Permanent atrial fibrillation | 148.21 | |
| Not to be confused with persistent; assign only when permanent AF is document | | |
| Unspecified atrial fibrillation | 148.91 | |
| Unspecified atrial flutter | 148.92 | |
| Sick sinus syndrome | 149.5 | |
| Left ventricular failure, unspecified | 150.1 | |

PRV20437-2501 As of 10/22/2024

Unspecified systolic (congestive) heart failure

Acute systolic (congestive) heart failure

150.20

150.21

F50.01

F50.02

F50.2

| Chronic systolic (congestive) heart failure Acute on chronic systolic (congestive) heart failure | 150.22 |
|---|---------|
| Acute on chronic systolic (congestive) heart failure | 150.00 |
| riodio ori oriio dyotolio (corigodiivo) riodic idiidio | 150.23 |
| Unspecified diastolic (congestive) heart failure | 150.30 |
| Acute diastolic (congestive) heart failure | 150.31 |
| Chronic diastolic (congestive) heart failure | 150.32 |
| Acute on chronic diastolic (congestive) heart failure | 150.33 |
| Unspecified combined systolic (congestive) and diastolic (congestive) heart failure | 150.40 |
| Acute combined systolic (congestive) and diastolic (congestive) heart failure | 150.41 |
| Chronic combined systolic (congestive) and diastolic (congestive) heart failure | 150.42 |
| Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure | 150.43 |
| Acute right heart failure | 150.811 |
| Acute on chronic right heart failure | 150.813 |
| End stage heart failure | 150.84 |
| Use additional code to identify the type of heart failure as systolic, diastolic or combined, if known (I50.2-I50.43). | |
| Heart failure, unspecified | 150.9 |
| Atherosclerosis of native arteries of extremities with rest pain, right leg | 170.221 |
| Atherosclerosis of native arteries of extremities with rest pain, left leg | 170.222 |
| Atherosclerosis of native arteries of extremities with rest pain, bilateral legs | 170.223 |
| Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, right leg | 170.321 |
| Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, left leg | 170.322 |
| Diagona of the Despiretory Cretory | CODE |
| Diseases of the Respiratory System | CODE |

| Diseases of the Respiratory System | CODE | |
|--|--------------|--|
| Unspecified chronic bronchitis | J42 | |
| Emphysema, unspecified | J43.9 | |
| Chronic obstructive pulmonary disease with (acute) lower respiratory infection | J44.0 | |
| Use additional code to identify the infection. | | |
| Chronic obstructive pulmonary disease with (acute) exacerbation | J44.1 | |
| If COPD with acute exacerbation and COPD with lower respiratory infection are present: document and code both. | | |
| Chronic obstructive pulmonary disease, unspecified | J44.9 | |
| Severe persistent asthma, uncomplicated | J45.50 | |
| Severe persistent asthma with (acute) exacerbation | J45.51 | |
| Severe persistent asthma with status asthmaticus | J45.52 | |
| Respiratory failure | J96.1-J96.92 | |
| Discours of the Discotive System | CODE | |

| Diseases of the Digestive System | CODE |
|----------------------------------|--------|
| Crohn's Disease, unspecified | K50.9- |
| Ulcerative colitis, unspecified | K51.9- |

| Diseases of the Musculoskeletal System and Connective Tissue | CODE | |
|--|-------------|--|
| Rheumatoid arthritis, unspecified | M06.9 | |
| Rheumatoid arthritis: If known, code specifically regarding with or without rheumatoid factor. Also code specific site as documented. | M05-M06.8A | |
| Systemic lupus erythematosus, unspecified | M32.9 | |
| Systemic lupus erythematosus (SLE): Documentation must confirm and specify Lupus as systemic to assign a code from category M32. | M32.0-M32.9 | |

For more coding and documentation tips, visit AlabamaBlue.com/Providers/CodingCorner.

| | Genitourinary Conditions | CODE |
|--|---|----------------|
| Two separate eGFR values are required to change the CKD stage. When both kidney failure and CKD are present, code both conditions. | | hen both acute |
| | Chronic kidney disease, stage 3 unspecified | N18.30 |
| | Chronic kidney disease, stage 3a | N18.31 |
| | Chronic kidney disease, stage 3b | N18.32 |
| | Chronic kidney disease, stage 4 (severe) | N18.4 |
| | Chronic kidney disease, stage 5 | N18.5 |
| | End stage renal disease | N18.6 |

| Factors Influencing Health Status | CODE |
|-----------------------------------|--------------|
| Body mass index (BMI) | Z68.1-Z68.45 |

BMI: Reported BMI >40 requires a weight diagnosis to be documented and coded, such as "Morbid Obesity" to rise to an HCC.

| Long-term (current) use of insulin | Z79.4 |
|------------------------------------|-------|
| | |

If the patient is treated with both oral hypoglycemic drugs and insulin, code both Z79.84 and Z79.4.

| Personal history of malignant neoplasm (code specific history of | Z85 |
|--|-----|
| cancer code) | |

Differentiating between "active" and "history of" cancer: When determining active versus history of cancer, look for active treatment, such as chemotherapy, radiation, hormone therapy, watchful waiting, or patient refusal of treatment. Remember, when coding active cancer, document specific primary sites and metastasis (secondary), if present.

| Personal history of transient ischemic attack (TIA)/cerebrovascular | Z86.73 |
|---|--------|
| accident (CVA) without residual effects | |

TIA/CVA: Use history of codes unless patient is having active symptoms of TIA or CVA during the visit. If the patient has residual of a CVA, code the residual symptoms or sequelae to the highest level of specificity I69.3- (i.e., monoplegia, hemiplegia, hemiplegia, hemiparesis, etc.).

| Acquired absence of toe(s), foot and ankle | Z89.4-Z89.449 |
|--|---------------|
| Acquired absence of leg below knee | Z89.5-Z89.519 |
| Acquired absence of leg above knee | Z89.6-Z89.619 |
| Tracheostomy status | Z93.0 |
| Gastrostomy status | Z93.1 |
| lleostomy status | Z93.2 |
| Colostomy status | Z93.3 |
| Cystostomy status | Z93.5- Z93.59 |
| Artificial opening status, unspecified | Z93.9 |
| Heart transplant status | Z94.1 |
| Lung transplant status | Z94.2 |
| Heart and lung transplant status | Z94.3 |
| Liver transplant status | Z94.4 |
| Bone marrow transplant status | Z94.81 |
| Intestine transplant status | Z94.82 |
| Pancreas transplant status | Z94.83 |
| Stem cells transplant status | Z94.84 |
| Presence of fully implantable artificial heart | Z95.812 |

When your patient is immunocompromised while taking anti-rejection medication, consider documenting and coding D84.821 "Immunodeficiency due to drugs."

Dependence on renal dialysis

Z99.2

ICD-10 is the 10th revision of the International Statistical Classification of Diseases and Related Health Problems (ICD), a medical classification list by the World Health Organization (WHO).



Blue Advantage® PPO is provided by Blue Cross and Blue Shield of Alabama, an independent licensee of the Blue Cross and Blue Shield Association.